

MarinTrust Standard V2

By-product Fishery Assessment Plaice (Pleuronectes platessa) in ICES Subdivisions 24-32

MarinTrust Programme

Unit C, Printworks 22 Amelia Street London SE17 3BZ

E: standards@marin-trust.com

T: +44 2039 780 819



Table 1 Application details and summary of the assessment outcome

	Species:	Plaice (Pleuronectes platessa)	
	Geographical area:	FAO Fishing Area 27 Atlantic Northeast	
Fishery Under Assessment	Country of origin of the product:	Denmark	
	Stock:	Plaice in ICES Subdivisions 24-32 (Baltic Sea, excluding the Sound and Belt Seas)	
Date	14 March 2022		
Report Code	BP040		
Assessor	Geraldine Criquet		
Country of origin of the product - PASS	Denmark		
Country of origin of the product - FAIL	NA		

Application details and	d summary of the assess	sment outcome	
Company Name(s): FF	Skagen		
Country: Denmark			
Email address:		Applicant Code	e:
Certification Body Det	ails		
Name of Certification	Body:	Global Trust C	ertification
Assessor	Peer Reviewer	Assessment Days	Initial/Surveillance/ Re-approval
Geraldine Criquet	Conor Donnelly	0.5	Surveillance 1
Assessment Period	To March 2022		

Scope Details	
Main Species	Plaice (Pleuronectes platessa)
Stock	Plaice in ICES Subdivisions 24-32 (Baltic Sea, excluding the Sound and Belt Seas)
Fishery Location	FAO Fishing Area 27 Atlantic Northeast
Management Authority (Country/ State)	EU & Denmark
Gear Type(s)	Active and passive gears
Outcome of Assessment	
Peer Review Evaluation	Agree with recommendation
Recommendation	APPROVED

Table 2. Assessment Determination

Assessment Determination

If any species is categorised as Endangered or Critically Endangered on IUCN's Red List, or if it appears in the CITES appendices, it cannot be approved for use as Marin Trust raw material. Plaice (*Pleuronectes platessa*) is neither listed as Endangered or Critically Endangered on IUCN's Red List, nor listed in CITES appendices; therefore, plaice is eligible for approval for use as Marin Trust by-product raw material.

The EU multiannual plan for the Baltic Sea applies to bycatches of this stock taken when fishing for the target stocks described in the plan. Refence points are defined. The stock is classified as Category C.

Fishery removals of the stock are considered in the stock assessment processes so the stock **PASSES** Clause C1.1.

In the most recent stock assessment, the stock is considered to have a biomass above the limit reference point, the stocks **PASSES** Clause C1.2.

Therefore, plaice stock in ICES Subdivisions 24-32 is **APPROVED** for the production of fishmeal and fish oil under the current Marin Trust v 2.0 by-products.

Fishery Assessment Peer Review Comments

The assessor correctly classified plaice (*Pleuronectes platessa*) in ICES Subdivisions 24-32 (Baltic Sea, excluding the Sound and Belt Seas) as category C, this stock is managed, and reference points are defined.

Fishery removals are considered in the stock assessment process. The most recent stock assessment shows that the stock is above B_{lim}. Therefore, the stock is considered to have a biomass above the limit reference point.

Plaice (*Pleuronectes platessa*) in ICES Subdivisions 24-32 (Baltic Sea, excluding the Sound and Belt Seas) passes both Clauses C1.1 and C1.2 and is therefore approved under the Marin Trust Standard v.2.

lotes for On-site Auditor
IA



Species Categorisation

NB: If any species is categorised as Endangered or Critically Endangered on the IUCN Red List, or if it appears in CITES Appendix 1, it **cannot** be approved for use as an MarinTrust raw material.

IUCN Red list Category

By-product material from a species listed by IUCN (the International Union for Conservation of Nature) under the Red List for the following categories shall immediately fail the assessment;

- EXTINCT (E) AND EXTINCT IN THE WILD (EW)
- CRITICALLY ENDANGERED (CR) facing an extremely high risk of extinction in the wild.
- ENDANGERED (EN) facing a very high risk of extinction in the wild.

By-product material may be used from the following categories provided that all clauses in the MarinTrust standard are passed.

- VULNERABLE (VU) facing a high risk of extinction in the wild.
- NEAR THREATENED (NT) does not qualify for above now, but is close or is likely to qualify for, a threatened category in the near future.
- LEAST CONCERN (LC) Widespread and abundant.
- DATA DEFICIENT (DD) and NOT EVALUATED (NE)

Table 3 Species Categorisation Table

Common name	Latin name	Stock	Management	Category	IUCN Red List Category ¹	CITES Appendix 1 ²
Plaice	Pleuronectes platessa	Plaice in ICES Subdivisions 24-32 (Baltic Sea, excluding the Sound and Belt Seas)		С	LC	No

¹ https://www.iucnredlist.org/

² https://cites.org/eng/app/appendices.php

CATEGORY C SPECIES

In a by-product assessment, Category C species are those which are subject to a species-specific management regime and are usually targeted species in fisheries for human consumption.

Clause C1 should be completed for each Category C species. If there are no Category C species in the fishery under assessment, this section can be deleted. Where a species fails this Clause, it should be assessed as a Category D species instead.

Spe	cies	Name	Plaice (Pleuronectes platessa)	
C1	Catego	ory C Stock Sta	atus - Minimum Requirements	
CI	C1.1	-	ovals of the species in the fishery under assessment are included in the stock assessment are considered by scientific authorities to be negligible.	Yes
	C1.2	The species i	is considered, in its most recent stock assessment, to have a biomass above the limit oint (or proxy), OR removals by the fishery under assessment are considered by scientific to be negligible.	Yes
		ı	Clause outcome:	DASS

C1.1 Fishery removals of the species in the fishery under assessment are included in the stock assessment process, OR are considered by scientific authorities to be negligible.

The stock assessment type is an age-based analytical assessment SAM, input data include commercial catches. Plaice catches (landings and discards) for the 2002-2020 period are shown in Figure 1. Therefore, the stock **PASSES** Clause C1.1.




Figure 1. Plaice in subdivision 24-32. Catches for the 2002-2020 period.

C1.2 The species is considered, in its most recent stock assessment, to have a biomass above the limit reference point (or proxy), OR removals by the fishery under assessment are considered by scientific authorities to be negligible.

As per the most recent ICES advice, the spawning-stock size is above the proxy MSY B_{trigger} (Figure 2). Therefore, the assessor determines that, the stock is considered to have a biomass above the limit reference point, it **PASSES** Clause C1.2.






Figure 2. Plaice in subdivisions 24-32. SPiCT model results used for the evaluation of the stock (proxy MSY B_{trigger} is represented by the horizontal line). The dots represent indices from BITS-Q1 and Q4.

References

ICES. 2021. Plaice (*Pleuronectes platessa*) in subdivisions 24-32 (Baltic Sea, excluding the Sound and Belt Seas). In Report of the ICES Advisory Committee, 2021. ICES Advice 2021, ple.27.24-32, https://doi.org/10.17895/ices.advice.7818.

Freyhof, J. 2014. *Pleuronectes platessa*. *The IUCN Red List of Threatened Species* 2014: e.T135690A50018800. https://dx.doi.org/10.2305/IUCN.UK.2014-1.RLTS.T135690A50018800.en. Accessed on 14 March 2022. https://www.iucnredlist.org/species/135690/50018800

Links

-	
MarinTrust Standard clause	1.3.2.2
FAO CCRF	7.5.3
GSSI	D.3.04, D5.01



CATEGORY D SPECIES

Category D species are those which are not subject to a species-specific management regime. In the case of mixed trawl fisheries, Category D species may make up the majority of landings. The comparative lack of scientific information on the status of the population of the species means that a risk-assessment style approach must be taken.

D1	Species Name		
	Productivity Attribute	e Value	Score
	Average age at maturity (years)		
	Average maximum age (years)		
	Fecundity (eggs/spawning)		
	Average maximum size (cm)		
	Average size at maturity (cm)		
	Reproductive strategy		
	Mean trophic level		
		Average Productivity Score	
	Susceptibility Attribut	te Value	Score
	Overlap of adult species range with fisher	ry	
	Distribution		
	Habitat		
	Depth range		
	Selectivity		
	Post-capture mortality		
		Average Susceptibility Score	
		PSA Risk Rating (From Table D3)	
		Compliance rating	
Refere	nces		
Standa	urd clauses 1.3.2.2		



Table D2 - Productivity / Susceptibility attributes and scores.

Productivity attributes	Low productivity/ High risk	Medium productivity/ Medium risk	High productivity Low risk	
	Score 3	Score 2	Score 1	
Average age at maturity (years)	>4	2 to 4	<2	
Average maximum age (years)	>30	10 to 30	<10	
Fecundity (eggs/spawning)	<1 000	1 000 to 10 000	>10 000	
Average maximum size (cm)	>150	60 to 150	<60	
Average size at maturity (cm)	>150	30 to 150	<30	
Reproductive strategy	Live bearer, mouth brooder or significant parental investment	Demersal spawner "berried"	Broadcast spawner	
Mean trophic level	>3.25	2.5-3.25	<2.5	

Susceptibility at	tribut	es	High susceptibility/ High risk	Medium susceptibility/ Medium risk	Low susceptibility/ Low risk	
			Score 3	Score 2	Score 1	
Availability	i	Overlap of adult species range with fishery	>50% of stock occurs in the area fished	Between 25% and 50% of the stock occurs in the area fished	<25% of stock occurs in the area fished	
	2) [Distribution	Only in the country/ fishery	Limited range in the region	Throughout region/ global distribution	
Encounterability	1) 1	Habitat	Habitat preference of species make it highly likely to encounter trawl gear (e.g. demersal, muddy/sandy bottom)	Habitat preference of species make it moderately likely to encounter trawl gear (e.g. rocky bottom/reefs)	Depth or distribution of species make it unlikely to encounter trawl gear (e.g. epi-pelagic or meso-pelagic)	
	2) [Depth range	High overlap with trawl fishing gear (20 to 60 m depth)	Medium overlap with trawl fishing gear (10 to 20 m depth)	Low overlap with trawl fishing gear (0 to 10 m, >70 m depth)	
Selectivity			Species >2 times mesh size or up to 4 m length	Species 1 to 2 times mesh size or 4 to 5 m length	Species <mesh or<br="" size="">>5 m length</mesh>	
Post capture mortality			Most dead or retained Trawl tow >3 hours	Alive after net hauled Trawl tow 0.5 to 3 hours	Released alive Trawl tow <0.5 hours	

Note: Availability 2 is only used when there is no information for Availability 1; the most conservative score between Encounterability 1 and 2 is used.



D3		Average Susceptibility Score			
		1 - 1.75	1.76 - 2.24	2.25 - 3	
Average Productivity	1 - 1.75	PASS	PASS	PASS	
Score	1.76 - 2.24	PASS	PASS	TABLE D4	
	2.25 - 3	PASS	TABLE D4	TABLE D4	

D4	Spe	cies Name				
	Impacts On Species Categorised as Vulnerable by D1-D3 - Minimum Requirements					
	D4.1	1 '	of the fishery on this species are considered during the management e measures are taken to minimise these impacts.			
	D4.2	There is no substantial species.	evidence that the fishery has a significant negative impact on the			
			Outcome:			
	-	ential impacts of the fis easures are taken to mini	hery on this species are considered during the management process, a mise these impacts.	nd		
reasor	nable me	easures are taken to mini		nd		
reasor	nable me	easures are taken to mini	mise these impacts.	nd		
reasor D4.2 T	nable me	easures are taken to mini	mise these impacts.	nd		
D4.2 T Refere	nable mo	easures are taken to mini	mise these impacts.	nd		
D4.2 T Refere	There is rences	easures are taken to mini no substantial evidence t	mise these impacts. nat the fishery has a significant negative impact on the species.	nd		