



## MarinTrust Standard V2

# By-product Fishery Assessment Report Template

**MarinTrust Programme**

Unit C, Printworks

22 Amelia Street

London

SE17 3BZ

E: [standards@marin-trust.com](mailto:standards@marin-trust.com)

T: +44 2039 780 819

**Table 1 Application details and summary of the assessment outcome**

Fishery Under Assessment	Species:	Norway Silvery cod / Silvery pout ( <i>Gadiculus argenteus thori</i> )
	Geographical area:	FAO Area 27 Atlantic, Northeast
	Country of origin of the product:	Norway
	Stock:	Norwegian silver cod stock
Date	19/10/2021	
Report Code	BP218	
Assessor	Virginia Polonio	
Country of origin of the product - PASS	Norway	
Country of origin of the product - FAIL	NA	

Application details and summary of the assessment outcome			
Name:			
Address:			
Country: Norway		Zip:	
Tel. No.:		Fax. No.:	
Email address:		Applicant Code:	
Key Contact:		Title:	
Certification Body Details			
Name of Certification Body:		Global Trust Certification	
Assessor	Peer Reviewer	Assessment Days	Initial/Surveillance/ Re-approval
Virginia Polonio	Geraldine Criquet	0.5	Re-approval
Assessment Period	To October 2021		

Scope Details	
Main Species	Norway Silvery cod / Silvery pout ( <i>Gadiculus argenteus thori</i> )
Stock	Norwegian silver cod stock
Fishery Location	FAO Area 27 Atlantic, Northeast
Management Authority (Country/ State)	Norway Department of Fisheries
Gear Type(s)	Bottom trawls
Outcome of Assessment	
Peer Review Evaluation	Agree with the assessor's recommendation of approval
Recommendation	<b>APPROVED</b>

## Table 2. Assessment Determination

Assessment Determination
<p>If any species is categorised as Endangered or Critically Endangered on the IUCN Red List, or if it appears in the CITES appendices, it cannot be approved for use as Marin Trust raw material. Norway Silvery cod / Silvery pout (<i>Gadiculus argenteus thori</i>) does not appear as Endangered or Critically Endangered on the IUCN Red List, nor does it appear in the CITES appendices; therefore, Silvery cod is eligible for approval for use as Marin Trust raw material.</p> <p>The species is not subject to research and specific management measures are not in place. Therefore, following Marin Trust criteria, the stock is classified as Category D.</p> <p>Due to the lack of scientific information on the status of the stock, a risk-assessment style approach was taken. The fishery was assessed using the risk-based Productivity, Susceptibility Analysis (PSA) approach as per Marin Trust v 2.0 procedures for Category D species.</p> <p>Table D1 (PSA) has achieved a PASS therefore, the average for the PSA risk rating results in the species passing D1.</p> <p>Norway Silvery cod / Silvery pout (<i>Gadiculus argenteus thori</i>) is APPROVED for the production of fishmeal and fish oil under the Marintrust v 2.0 by-products standard.</p>
Fishery Assessment Peer Review Comments
<p>The assessor correctly classified the Norway silver cod stock as category D, reference points are not defined to assess status of the stock relative to.</p> <p>With an average productivity score of 1.33 and an average susceptibility score of 2.5, the Norway silver cod PASSES Clause D1 in accordance with Table D3.</p> <p>Therefore, the Norway silver cod stock should be approved.</p>
Notes for On-site Auditor

## Species Categorisation

**NB:** If any species is categorised as Endangered or Critically Endangered on the IUCN Red List, or if it appears in CITES Appendix 1, it **cannot** be approved for use as an MARINTRUST raw material.

### IUCN Red list Category

By-product material from a species listed by IUCN (the International Union for Conservation of Nature) under the Red List for the following categories shall immediately fail the assessment;

- EXTINCT (E) AND EXTINCT IN THE WILD (EW)
- CRITICALLY ENDANGERED (CR) facing an extremely high risk of extinction in the wild.
- ENDANGERED (EN) facing a very high risk of extinction in the wild.

By-product material may be used from the following categories provided that all clauses in the MarinTrust standard are passed.

- VULNERABLE (VU) facing a high risk of extinction in the wild.
- NEAR THREATENED (NT) does not qualify for above now, but is close or is likely to qualify for, a threatened category in the near future.
- LEAST CONCERN (LC) Widespread and abundant.
- DATA DEFICIENT (DD) and NOT EVALUATED (NE)

## Table 3 Species Categorisation Table

Common name	Latin name	Stock	Management	Category	IUCN Red List Category <sup>1</sup>	CITES Appendix 1 <sup>2</sup>
Norway Silvery cod / Silvery pout	<i>Gadiculus argenteus thori</i>	Norway silver cod	Norway	D	LC	No

<sup>1</sup> <https://www.iucnredlist.org/>

<sup>2</sup> <https://cites.org/eng/app/appendices.php>

## CATEGORY D SPECIES


Category D species are those which make up less than 5% of landings and are not subject to a species-specific management regime. In the case of mixed trawl fisheries, Category D species may make up the majority of landings. The comparative lack of scientific information on the status of the population of the species means that a risk-assessment style approach must be taken.

<b>D1</b>	<b>Species Name</b>	Norway Silvery cod / Silvery pout <i>Gadiculus argenteus thori</i>		
	<b>Productivity Attribute</b>		<b>Value</b>	<b>Score</b>
	Average age at maturity (years)		1	1
	Average maximum age (years)		3	1
	Fecundity (eggs/spawning)		No value	Not scored
	Average maximum size (cm)		15.00	1
	Average size at maturity (cm)		12.0	1
	Reproductive strategy		nonguarders: open water/substratum egg scatterers	1
	Mean trophic level		3.5	3
	<b>Average Productivity Score</b>			<b>1.33</b>
	<b>Susceptibility Attribute</b>		<b>Value</b>	<b>Score</b>
	Overlap of adult species range with fishery		25-50%	2
	Distribution		Not scored	-
	Habitat		Pelagic, muddy bottoms	2
	Depth range		100-1000m	1
	Selectivity		Up to 4m length	3
	Post-capture mortality		Retained	3
<b>Average Susceptibility Score</b>			<b>2.5</b>	
<b>PSA Risk Rating (From Table D3)</b>			<b>PASS</b>	
<b>Compliance rating</b>			<b>PASS</b>	

### References

Productivity attributes:

Fishbase. <https://www.fishbase.de/Summary/SpeciesSummary.php?ID=25&AT=silvery+cod>



\*Figure 1. Distribution maps for *Gadiculus thori*, with modelled year 2050 native range map based on IPCC RCP8.5 emissions scenario. [www.aquamaps.org](http://www.aquamaps.org), version 10/2019.

Scarponi, P., G. Coro, and P. Pagano. A collection of Aquamaps native layers in NetCDF format. Data in brief 17 (2018): 292-296.

Standard clauses 1.3.2.2

Table D2 - Productivity / Susceptibility attributes and scores.

Productivity attributes	Low productivity/ High risk	Medium productivity/ Medium risk	High productivity/ Low risk
	Score 3	Score 2	Score 1
Average age at maturity (years)	>4	2 to 4	<2
Average maximum age (years)	>30	10 to 30	<10
Fecundity (eggs/spawning)	<1 000	1 000 to 10 000	>10 000
Average maximum size (cm)	>150	60 to 150	<60
Average size at maturity (cm)	>150	30 to 150	<30
Reproductive strategy	Live bearer, mouth brooder or significant parental investment	Demersal spawner "berried"	Broadcast spawner
Mean trophic level	>3.25	2.5–3.25	<2.5

Susceptibility attributes		High susceptibility/ High risk	Medium susceptibility/ Medium risk	Low susceptibility/ Low risk
		Score 3	Score 2	Score 1
Availability	1) Overlap of adult species range with fishery	>50% of stock occurs in the area fished	Between 25% and 50% of the stock occurs in the area fished	<25% of stock occurs in the area fished
	2) Distribution	Only in the country/ fishery	Limited range in the region	Throughout region/ global distribution
Encounterability	1) Habitat	Habitat preference of species make it highly likely to encounter trawl gear (e.g. demersal, muddy/sandy bottom)	Habitat preference of species make it moderately likely to encounter trawl gear (e.g. rocky bottom/reefs)	Depth or distribution of species make it unlikely to encounter trawl gear (e.g. epi-pelagic or meso-pelagic)
	2) Depth range	High overlap with trawl fishing gear (20 to 60 m depth)	Medium overlap with trawl fishing gear (10 to 20 m depth)	Low overlap with trawl fishing gear (0 to 10 m, >70 m depth)
Selectivity		Species >2 times mesh size or up to 4 m length	Species 1 to 2 times mesh size or 4 to 5 m length	Species <mesh size or >5 m length
Post capture mortality		Most dead or retained Trawl tow >3 hours	Alive after net hauled Trawl tow 0.5 to 3 hours	Released alive Trawl tow <0.5 hours

**Note:** Availability 2 is only used when there is no information for Availability 1; the most conservative score between Encounterability 1 and 2 is used.

D3		Average Susceptibility Score		
		1 - 1.75	1.76 - 2.24	2.25 - 3
Average Productivity Score	1 - 1.75	PASS	PASS	PASS
	1.76 - 2.24	PASS	PASS	TABLE D4
	2.25 - 3	PASS	TABLE D4	TABLE D4

D4 Species Name			
<b>Impacts On Species Categorised as Vulnerable by D1-D3 - Minimum Requirements</b>			
D4.1	The potential impacts of the fishery on this species are considered during the management process, and reasonable measures are taken to minimise these impacts.		
D4.2	There is no substantial evidence that the fishery has a significant negative impact on the species.		
<b>Outcome:</b>			
<b>Evidence</b>			
D4.1: The potential impacts of the fishery on this species are considered during the management process, and reasonable measures are taken to minimise these impacts.			
D4.2 There is no substantial evidence that the fishery has a significant negative impact on the species.			
<b>References</b>			
<b>Links</b>			
MARINTRUST Standard clause	1.3.2.2, 4.1.4		
FAO CCRF	7.5.1		
GSSI	D.5.01		