

# **IFFO RS**Global Standard for Responsible Supply of Marine Ingredients



#### **IFFO RS Limited**

T: +44 (0) 2030 539 195 E: Standards@iffors.com W: www.iffors.com

Unit C, Printworks | 22 Amelia Street London, SE17 3BZ | United Kingdom





Global Standard for Responsible Supply of Marine Ingredients Fishery Assessment Methodology and Template Report V2.0



# **IFFO RS**Global Standard for Responsible Supply of Marine Ingredients



Fishery Under Assessment	Black bream / black seabream  Spondyliosoma cantharus
Date	January 2019
Assessor	Conor Donnelly

Application details and summary of the assessment outcome				
Name: Copalis Indus	trie			
Address:				
Country:		Zip:		
Tel. No.:		Fax. No.:		
Email address:		Applicant Code		
Key Contact:		Title:		
Certification Body Details				
Name of Certification	Body:			
Assessor Name	Peer Reviewer	Assessment Days	Initial/Surveillance/Re approval	Whole fish/ By- product
Conor Donnelly Jim Daly		1	Surveillance	By-product
Assessment Period	2018			

Scope Details	
Management Authority (Country/State)	EU
Main Species	Black bream/ Black seabream Spondyliosoma cantharus
Fishery Location	EU waters
Gear Type(s)	Benthic and demersal trawls
Outcome of Assessment	
Overall Outcome	Pass
Clauses Failed	None
Peer Review Evaluation	Pass
Recommendation	Approval

#### **Assessment Determination**

The comparative lack of scientific information on the status of the population in the assessment area means that a risk-assessment style approach must be taken. The species was assessed using the risk-based Productivity, Susceptibility Analysis (PSA) strategy as per IFFO-RS v 2.0 Standard for Category D species. The species passes this risk-based approach.

The IUCN has categorised *Spondyliosoma cantharus* as a species of least concern (European assessment, last assessed 27 May 2013), it is not listed on the current CITES appendices.

Consequently, the assessment team recommends approving this byproduct material against the IFFO RS  $\nu$  2.0 Standard for the production of fishmeal and fish oil.

#### **Peer Review Comments**

Agree with assessment.

#### **Notes for On-site Auditor**

Species-Specific Results

Category	Species	% landings	Outcome (Pass/Fail)
			A1
Cotogory			A2
Category A			A3
			A4
Category B			
Category C			
Category D	Black bream Spondyliosoma cantharus	NA	Pass

[List all Category A and B species. List approximate total % age of landings which are Category C and D species; these do not need to be individually named here]

#### HOW TO COMPLETE THIS ASSESSMENT REPORT

This assessment template uses a modular approach to assessing fisheries against the IFFO RS standard.

#### Whole Fish

The process for completing the template for a **whole fish** assessment is as follows:

- 1. ALL ASSESSMENTS: Complete the Species Characterisation table, to determine which categories of species are present in the fishery.
- 2. ALL ASSESSMENTS: Complete clauses M1, M2, M3: Management.
- 3. IF THERE ARE CATEGORY A SPECIES IN THE FISHERY: Complete clauses A1, A2, A3, A4 for **each** Category A species.
- 4. IF THERE ARE CATEGORY B SPECIES IN THE FISHERY: Complete the Section B risk assessment for **each** Category B species.
- 5. IF THERE ARE CATEGORY C SPECIES IN THE FISHERY: Complete clause C1 for **each** Category C species.
- 6. IF THERE ARE CATEGORY D SPECIES IN THE FISHERY: Complete Section D.
- 7. ALL ASSESSMENTS: Complete clauses F1, F2, F3: Further Impacts.

A fishery must score a pass in **all applicable clauses** before approval may be recommended. To achieve a pass in a clause, the fishery/species must meet **all** of the minimum requirements.

#### By-products

The process for completing the template for **by-product raw material** is as follows:

- 1. ALL ASSESSMENTS: Complete the Species Characterisation table with the names of the by-product species and stocks under assessment. The '% landings' column can be left empty; all by-products are considered as Category C and D.
- 2. IF THERE ARE CATEGORY C BYPRODUCTS UNDER ASSESSMENT: Complete clause C1 for **each** Category C by-product.
- 3. IF THERE ARE CATEGORY D BYPRODUCTS UNDER ASSESSMENT: Complete Section D.
- 4. ALL OTHER SECTIONS CAN BE DELETED. Clauses M1 M3, F1 F3, and Sections A and B do not need to be completed for a by-product assessment.

By-product approval is awarded on a species-by-species basis. Each by-product species scoring a pass under the appropriate section may be approved against the IFFO RS Standard.

Version No.: 2.0 Date: July 2017 Page 5

#### SPECIES CATEGORISATION

The following table should be completed as fully as the available information permits. Any species representing more than 0.1% of the annual catch should be listed, along with an estimate of the proportion of the catch each species represents. The species should then be divided into Type 1 and Type 2 as follows:

- **Type 1 Species** can be considered the 'target' or 'main' species in the fishery. They make up the bulk of annual landings and are subjected to a detailed assessment.
- **Type 2 Species** can be considered the 'bycatch' or 'minor' species in the fishery. They make up a small proportion of the annual landings and are subjected to relatively high-level assessment.

Type 1 Species must represent 95% of the total annual catch. Type 2 Species may represent a maximum of 5% of the annual catch (see Appendix B).

Species which make up less than 0.1% of landings do not need to be listed (NOTE: ETP species are considered separately). The table should be extended if more space is needed. Discarded species should be included when known.

The 'stock' column should be used to differentiate when there are multiple biological or management stocks of one species captured by the fishery. The 'management' column should be used to indicate whether there is an adequate management regime specifically aimed at the individual species/stock. In some cases it will be immediately clear whether there is a species-specific management regime in place (for example, if there is an annual TAC). In less clear circumstances, the rule of thumb should be that if the species meets the minimum requirements of clauses A1-A4, an adequate species-specific management regime is in place.

NOTE: If any species is categorised as Endangered or Critically Endangered on the IUCN Red List, or if it appears in the CITES appendices, it **cannot** be approved for use as an IFFO RS raw material. This applied to whole fish as well as by-products.

## **TYPE 1 SPECIES (Representing 95% of the catch or more)**

Category A: Species-specific management regime in place.

Category B: No species-specific management regime in place.

#### **TYPE 2 SPECIES (Representing 5% OF THE CATCH OR LESS)**

Category C: Species-specific management regime in place.

Category D: No species-specific management regime in place.

Common name	Latin name	Stock	% of landings	Management	Category
Black bream	Spondyliosoma cantharus	EU waters		EU	D

Version No.: 2.0 Date: July 2017 Page 6

#### **CATEGORY D SPECIES**

In a whole fish assessment, Category D species are those which make up less than 5% of landings and are not subject to a species-specific management regime. In the case of mixed trawl fisheries, Category D species may make up the majority of landings. In a by-product assessment, Category D species are those which are not subject to a species-specific management regime. In both cases, the comparative lack of scientific information on the status of the population of the species means that a risk-assessment style approach must be taken.

The process for assessing Category D species involves the use of a Productivity-Susceptibility Analysis (PSA) to further subdivide the species into 'Critical Risk', 'Major Risk' and 'Minor Risk' groups. If there are no Category D species in the fishery under assessment, this section can be deleted.

Productivity and susceptibility ratings are calculated using a process derived from the APFIC document "Regional Guidelines for the Management of Tropical Trawl Fisheries, which in turn was derived from papers by Patrick *et al* (2009) and Hobday *et al* (2007). Table D1 should be completed for each Category D species as follows:


- Firstly, the best available information should be used to fill in values for each productivity and susceptibility attribute.
- Table D2 should be used to convert each attribute value into a score between 1 and 3.
- The average score for productivity attributes and the average for susceptibility attributes should be calculated.
- Table D3 should be used to determine whether the species is required to meet the requirements of Table D4. A species which does not need to meet the requirements of D4 is automatically awarded a pass.
- Table D4 should be used to assess those species indicated by Table D3 to determine a pass/fail rating.
- Any Category D species which has been categorised by the IUCN Red List as Endangered or Critically Endangered, or which appears in the CITES appendices, automatically results in a fail.

<b>Species Name:</b>	Black seabream Sp	pondyliosoma cantharus	
<b>Productivity Attribute</b>		Value	Score
Average age at maturity	(years)	2.8	2
Average maximum age	(years)	10.2	2
Fecundity (eggs/spawning	ng)	31,670-554,070	1
Average maximum size	(cm)	37.02	1
Average size at maturity	(cm)	20.8	1
Reproductive strategy		Non-guarders; brood hiders.	2
Mean trophic level		3.3	3
		Average Productivity Score	1.71
Susceptibility Attribut	e	Value	Score
Overlap of adult species	range with fishery	50%	3
Distribution		Not scored when overlap scored	n/a
Habitat		Benthopelagic	3
Depth range		5-300m	3
Selectivity		Up to 4m length	3
Post-capture mortality		Most dead or retained	3
		Average Susceptibility Score	3
		1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
		PSA Risk Rating (From Table D3)	Pass

#### **References:**

## R1 Overlap and distribution attribute:

Version No.: 2.0 Date: July 2017 Page 7



Native distribution map for *Spondyliosoma cantharus*. (Source: <a href="https://www.aquamaps.org/receive.php?type\_of\_map=regular#">https://www.aquamaps.org/receive.php?type\_of\_map=regular#</a>)

#### Other attributes:

R2 Fishbase: https://www.fishbase.de/summary/1356

R3 IUCN Red List <a href="https://www.iucnredlist.org/species/170258/42450257">https://www.iucnredlist.org/species/170258/42450257</a>

R4 CITES Checklist:

http://checklist.cites.org/#/en/search/output\_layout=alphabetical&level\_of\_listing=0&show\_synonyms=1&show\_author=1&show\_english=1&show\_spanish=1&show\_french=1&scientific\_name=black+bream&pag\_e=1&per\_page=20

Standard clauses 1.3.2.2

Table D2 - Productivity / Susceptibility attributes and scores.

Productivity attributes	Low productivity/ High risk	Medium productivity/ Medium risk	High productivity/ Low risk	
	Score 3	Score 2	Score 1	
Average age at maturity (years)	>4	2 to 4	<2	
Average maximum age (years)	>30	10 to 30	<10	
Fecundity (eggs/spawning)	<1 000	1 000 to 10 000	>10 000	
Average maximum size (cm)	>150	60 to 150	<60	
Average size at maturity (cm)	>150	30 to 150	<30	
Reproductive strategy	Live bearer, mouth brooder or significant parental investment	Demersal spawner "berried"	Broadcast spawner	
Mean trophic level	>3.25	2.5-3.25	<2.5	

Susceptibility attributes		High susceptibility/ High risk	Medium susceptibility/ Medium risk	Low susceptibility/ Low risk	
			Score 3	Score 2	Score 1
Availability	1)	Overlap of adult species range with fishery	>50% of stock occurs in the area fished	Between 25% and 50% of the stock occurs in the area fished	<25% of stock occurs in the area fished
	2)	Distribution	Only in the country/ fishery	Limited range in the region	Throughout region/ global distribution
Encounterability	1)	Habitat	Habitat preference of species make it highly likely to encounter trawl gear (e.g. demersal, muddy/sandy bottom)	Habitat preference of species make it moderately likely to encounter trawl gear (e.g. rocky bottom/reefs)	Depth or distribution of species make it unlikely to encounter trawl gear (e.g. epi-pelagic or meso-pelagic)
	2)	Depth range	High overlap with trawl fishing gear (20 to 60 m depth)	Medium overlap with trawl fishing gear (10 to 20 m depth)	Low overlap with trawl fishing gear (0 to 10 m, >70 m depth)
Selectivity			Species >2 times mesh size or up to 4 m length	Species 1 to 2 times mesh size or 4 to 5 m length	Species <mesh or<br="" size="">&gt;5 m length</mesh>
Post capture mortality			Most dead or retained Trawl tow >3 hours	Alive after net hauled Trawl tow 0.5 to 3 hours	Released alive Trawl tow <0.5 hours

**Note:** Availability 2 is only used when there is no information for Availability 1; the most conservative score between Encounterability 1 and 2 is used.

D3		Ave	erage Susceptibility So	core
D3		1.00 – 1.75	1.76 – 2.24	2.25 - 3.00
Average Productivity	1.00 - 1.75	PASS	PASS	PASS
Score	1.76 – 2.24	PASS	PASS	TABLE D4
	2.25 - 3.00	PASS	TABLE D4	TABLE D4

<b>D4</b>	Spec	cies Name	
	Impa	cts On Species Catego	orised as Vulnerable by D1-D3 - Minimum Requirements
	D4.1		cts of the fishery on this species are considered during the s, and reasonable measures are taken to minimise these impacts.
	D4.2		ial evidence that the fishery has a significant negative impact on
		the species.	Outcome:
Evide	nce		
Refer			
Stande	ard clau	ise 1.3.2.2	

# **SOCIAL CRITERION**

In addition to the scored criteria listed above, applicants must commit to ensuring that vessels operating in the fishery adhere to internationally recognised guidance on human rights. They must also commit to ensuring there is no use of enforced or unpaid labour in the fleet(s) operating upon the resource.