



RESPONSIBLE
SUPPLY

IFFO RS
Global Standard for Responsible Supply
of Marine Ingredients

IFFO RS Limited

T: +44 (0) 2030 539 195
E: Standards@iffors.com
W: www.iffors.com

Unit C, Printworks | 22 Amelia Street
London, SE17 3BZ | United Kingdom



**Global Standard for
Responsible Supply
of Marine Ingredients**
Fishery Assessment
Methodology and
Template Report V2.0



RESPONSIBLE
SUPPLY

IFFO RS

Global Standard for Responsible Supply
of Marine Ingredients



Fishery Under Assessment	Bullet tuna <i>Auxis rochei</i> Eastern Pacific Ocean
Date	October 2019
Assessor	Jim Daly

Application details and summary of the assessment outcome				
Name: Productos Pesqueros S.A Produpes				
Address:				
Country: Ecuador		Zip:		
Tel. No.:		Fax. No.:		
Email address:		Applicant Code		
Key Contact:		Title:		
Certification Body Details				
Name of Certification Body:		SAI Global Ltd		
Assessor Name	Peer Reviewer	Assessment Days	Initial/Surveillance/R e-approval	Whole fish/ By-product
Jim Daly	Conor Donnelly	0.5	Surveillance 2	By-product
Assessment Period	2019			

Scope Details	
Management Authority (Country/State)	Inter-American Tropical Tuna Commission (IATTC)
Main Species	Bullet tuna (<i>Auxis rochei</i>)
Fishery Location	FAO 87
Gear Type(s)	Purse seine longlines
Outcome of Assessment	
Overall Outcome	Pass
Clauses Failed	None
Peer Review Evaluation	Approve
Recommendation	Pass

Assessment Determination

As a highly-migratory species, effective management of Bullet tuna stocks is necessarily international. Bullet tuna fisheries in FAO87, the Eastern Pacific, fall under the jurisdiction of the Inter-American Tropical Tuna Commission (IATTC). The IATTC produces regular fishery status reports, the most recent of which was published in 2019, reporting on the 2018 tuna fisheries in the assessment area (IATTC, 2019).

Bullet tuna is considered less important than principle tuna species is considered together with similar species under 'Other tunas and tuna-like species. No species-specific management measures or stock assessments are available for this species IATTC website accessed 18.10.19)

The comparative lack of scientific information on the status of the population in the assessment area means that a risk-assessment style approach must be taken. The fishery was assessed using the risk-based Productivity, Susceptibility Analysis (PSA) as per IFFO-RS v 2.0 procedures for Category D species. The species has passed this risk-based assessment (**Table D3**).

The IUCN has categorised Bullet tuna *Auxis rochei* as a species of least concern; it is not currently listed on the CITES appendices of species most endangered of extinction (websites accessed 18.10.19).

Consequently, the assessment team recommends approving this by-product material against the IFFO RS v 2.0 Standard for the production of fishmeal and fish oil.

Peer Review Comments

Agree

Notes for On-site Auditor

Species-Specific Results

Category	Species	% landings	Outcome (Pass/Fail)
Category D	Bullet tuna <i>Auxis rochei</i>	N/A	Pass

[List all Category A and B species. List approximate total %age of landings which are Category C and D species; these do not need to be individually named here]

HOW TO COMPLETE THIS ASSESSMENT REPORT

This assessment template uses a modular approach to assessing fisheries against the IFFO RS standard.

Whole Fish

The process for completing the template for a **whole fish** assessment is as follows:

1. ALL ASSESSMENTS: Complete the Species Characterisation table, to determine which categories of species are present in the fishery.
2. ALL ASSESSMENTS: Complete clauses M1, M2, M3: Management.
3. IF THERE ARE CATEGORY A SPECIES IN THE FISHERY: Complete clauses A1, A2, A3, A4 for **each** Category A species.
4. IF THERE ARE CATEGORY B SPECIES IN THE FISHERY: Complete the Section B risk assessment for **each** Category B species.
5. IF THERE ARE CATEGORY C SPECIES IN THE FISHERY: Complete clause C1 for **each** Category C species.
6. IF THERE ARE CATEGORY D SPECIES IN THE FISHERY: Complete Section D.
7. ALL ASSESSMENTS: Complete clauses F1, F2, F3: Further Impacts.

A fishery must score a pass in **all applicable clauses** before approval may be recommended. To achieve a pass in a clause, the fishery/species must meet **all** of the minimum requirements.

By-products

The process for completing the template for **by-product raw material** is as follows:

1. ALL ASSESSMENTS: Complete the Species Characterisation table with the names of the by-product species and stocks under assessment. The '% landings' column can be left empty; all by-products are considered as Category C and D.
2. IF THERE ARE CATEGORY C BYPRODUCTS UNDER ASSESSMENT: Complete clause C1 for **each** Category C by-product.
3. IF THERE ARE CATEGORY D BYPRODUCTS UNDER ASSESSMENT: Complete Section D.
4. ALL OTHER SECTIONS CAN BE DELETED. Clauses M1 - M3, F1 - F3, and Sections A and B do not need to be completed for a by-product assessment.

By-product approval is awarded on a species-by-species basis. Each by-product species scoring a pass under the appropriate section may be approved against the IFFO RS Standard.

SPECIES CATEGORISATION

The following table should be completed as fully as the available information permits. Any species representing more than 0.1% of the annual catch should be listed, along with an estimate of the proportion of the catch each species represents. The species should then be divided into Type 1 and Type 2 as follows:

- **Type 1 Species** can be considered the 'target' or 'main' species in the fishery. They make up the bulk of annual landings and are subjected to a detailed assessment.
- **Type 2 Species** can be considered the 'bycatch' or 'minor' species in the fishery. They make up a small proportion of the annual landings and are subjected to relatively high-level assessment.

Type 1 Species must represent 95% of the total annual catch. Type 2 Species may represent a maximum of 5% of the annual catch (see Appendix B).

Species which make up less than 0.1% of landings do not need to be listed (NOTE: ETP species are considered separately). The table should be extended if more space is needed. Discarded species should be included when known.

The 'stock' column should be used to differentiate when there are multiple biological or management stocks of one species captured by the fishery. The 'management' column should be used to indicate whether there is an adequate management regime specifically aimed at the individual species/stock. In some cases, it will be immediately clear whether there is a species-specific management regime in place (for example, if there is an annual TAC). In less clear circumstances, the rule of thumb should be that if the species meets the minimum requirements of clauses A1-A4, an adequate species-specific management regime is in place.

NOTE: If any species is categorised as Endangered or Critically Endangered on the IUCN Red List, or if it appears in the CITES appendices, it **cannot** be approved for use as an IFFO RS raw material. This applied to whole fish as well as by-products.

TYPE 1 SPECIES (Representing 95% of the catch or more)

Category A: Species-specific management regime in place.

Category B: No species-specific management regime in place.

TYPE 2 SPECIES (Representing 5% OF THE CATCH OR LESS)

Category C: Species-specific management regime in place.

Category D: No species-specific management regime in place.

Common name	Latin name	Stock	% of landings	Management	Category
Bullet tuna	<i>Auxis rochei</i>	FAO 87	N/A	IATTC	D

CATEGORY D SPECIES

In a whole fish assessment, Category D species are those which make up less than 5% of landings and are not subject to a species-specific management regime. In the case of mixed trawl fisheries, Category D species may make up the majority of landings. In a by-product assessment, Category D species are those which are not subject to a species-specific management regime. In both cases, the comparative lack of scientific information on the status of the population of the species means that a risk-assessment style approach must be taken.

The process for assessing Category D species involves the use of a Productivity-Susceptibility Analysis (PSA) to further subdivide the species into 'Critical Risk', 'Major Risk' and 'Minor Risk' groups. If there are no Category D species in the fishery under assessment, this section can be deleted.

Productivity and susceptibility ratings are calculated using a process derived from the APFIC document "Regional Guidelines for the Management of Tropical Trawl Fisheries, which in turn was derived from papers by Patrick *et al* (2009) and Hobday *et al* (2007). Table D1 should be completed for each Category D species as follows:

- Firstly, the best available information should be used to fill in values for each productivity and susceptibility attribute.
- Table D2 should be used to convert each attribute value into a score between 1 and 3.
- The average score for productivity attributes and the average for susceptibility attributes should be calculated.
- Table D3 should be used to determine whether the species is required to meet the requirements of Table D4. A species which does not need to meet the requirements of D4 is automatically awarded a pass.
- Table D4 should be used to assess those species indicated by Table D3 to determine a pass/fail rating.
- Any Category D species which has been categorised by the IUCN Red List as Endangered or Critically Endangered, or which appears in the CITES appendices, automatically results in a fail.

D1	Species Name:	Bullet tuna <i>Auxis rochei</i>	
	Productivity Attribute	Value	Score
	Average age at maturity (years)	2	2
	Average maximum age (years)	5	1
	Fecundity (eggs/spawning)	31,000 – 103,000	1
	Average maximum size (cm)	50	1
	Average size at maturity (cm)	35	2
	Reproductive strategy	Broadcast spawner	1
	Mean trophic level	4.4	3
	Average		1.57
	Productivity Score		
	Susceptibility Attribute	Value	Score
	Overlap of adult species range with fishery	No information	
	Distribution	Throughout region	1
	Habitat	Pelagic-neritic	N/A
	Depth range	0-200m	3
	Selectivity	Up to 4m length	3
	Post-capture mortality	Most dead or retained	3
Average Susceptibility Score		2.5	
PSA Risk Rating (From Table D3)		Pass	
Compliance rating		Pass	

References R1-R5




Figure 1 Native distribution map for *Auxis rochei* (Source: Aquamaps) **R1**

R1 Fishbase Aquamaps

<https://www.fishbase.de/Summary/SpeciesSummary.php?ID=93&AT=BULLET+TUNA>

R2 IUCN Redlist: <https://www.iucnredlist.org/> Species Search Bullet tuna

R3 Fishbase <https://www.fishbase.de/Summary/SpeciesSummary.php?ID=93&AT=BULLET+TUNA>

R4 IATTC resolutions and recommendations. <https://www.iattc.org/ResolutionsActiveENG.htm>

R5 IATTC: Fishery Status Report No. 17, (2019). Tunas, billfishes and other pelagic species in the eastern Pacific Ocean in 2018: http://www.iattc.org/PDFFiles/FisheryStatusReports/English/No-17-2019_Tuna%20fishery,%20stocks,%20and%20ecosystem%20in%20the%20eastern%20Pacific%20Ocean%20in%202018.pdf

Standard clauses 1.3.2.2

Table D2 - Productivity / Susceptibility attributes and scores.

Productivity attributes	Low productivity/ High risk	Medium productivity/ Medium risk	High productivity/ Low risk
	Score 3	Score 2	Score 1
Average age at maturity (years)	>4	2 to 4	<2
Average maximum age (years)	>30	10 to 30	<10
Fecundity (eggs/spawning)	<1 000	1 000 to 10 000	>10 000
Average maximum size (cm)	>150	60 to 150	<60
Average size at maturity (cm)	>150	30 to 150	<30
Reproductive strategy	Live bearer, mouth brooder or significant parental investment	Demersal spawner "berried"	Broadcast spawner
Mean trophic level	>3.25	2.5–3.25	<2.5

Susceptibility attributes		High susceptibility/ High risk	Medium susceptibility/ Medium risk	Low susceptibility/ Low risk
		Score 3	Score 2	Score 1
Availability	1) Overlap of adult species range with fishery	>50% of stock occurs in the area fished	Between 25% and 50% of the stock occurs in the area fished	<25% of stock occurs in the area fished
	2) Distribution	Only in the country/fishery	Limited range in the region	Throughout region/global distribution
Encounterability	1) Habitat	Habitat preference of species make it highly likely to encounter trawl gear (e.g. demersal, muddy/sandy bottom)	Habitat preference of species make it moderately likely to encounter trawl gear (e.g. rocky bottom/reefs)	Depth or distribution of species make it unlikely to encounter trawl gear (e.g. epi-pelagic or meso-pelagic)
	2) Depth range	High overlap with trawl fishing gear (20 to 60 m depth)	Medium overlap with trawl fishing gear (10 to 20 m depth)	Low overlap with trawl fishing gear (0 to 10 m, >70 m depth)
Selectivity		Species >2 times mesh size or up to 4 m length	Species 1 to 2 times mesh size or 4 to 5 m length	Species <mesh size or >5 m length
Post capture mortality		Most dead or retained Trawl tow >3 hours	Alive after net hauled Trawl tow 0.5 to 3 hours	Released alive Trawl tow <0.5 hours

Note: Availability 2 is only used when there is no information for Availability 1; the most conservative score between Encounterability 1 and 2 is used.

D3		Average Susceptibility Score		
		1.00 – 1.75	1.76 – 2.24	2.25 – 3.00
Average Productivity Score	1.00 – 1.75	PASS	PASS	PASS
	1.76 – 2.24	PASS	PASS	TABLE D4
	2.25 – 3.00	PASS	TABLE D4	TABLE D4

D4	Species Name		
Impacts On Species Categorised as Vulnerable by D1-D3 - Minimum Requirements			
D4.1	The potential impacts of the fishery on this species are considered during the management process, and reasonable measures are taken to minimise these impacts.		
D4.2	There is no substantial evidence that the fishery has a significant negative impact on the species.		
Outcome:			
Evidence			
References			
<i>Standard clause 1.3.2.2</i>			

SOCIAL CRITERION

In addition to the scored criteria listed above, applicants must commit to ensuring that vessels operating in the fishery adhere to internationally recognised guidance on human rights. They must also commit to ensuring there is no use of enforced or unpaid labour in the fleet(s) operating upon the resource.

Appendix A - Determining Resilience Ratings

The assessment of Category B species described in this assessment report template utilises a resilience rating system suggested by the American Fisheries Society. This approach was chosen because it is also used by FishBase, and so the resilience ratings for many thousands of species are freely available online. As described by FishBase, the following is the process used to arrive at the resilience ratings:

"The American Fisheries Society (AFS) has suggested values for several biological parameters that allow classification of a fish population or species into categories of high, medium, low and very low resilience or productivity (Musick 1999). If no reliable estimate of r_m (see below) is available, the assignment is to the lowest category for which any of the available parameters fits. For each of these categories, AFS has suggested thresholds for decline over the longer of 10 years or three generations. If an observed decline measured in biomass or numbers of mature individuals exceeds the indicated threshold value, the population or species is considered vulnerable to extinction unless explicitly shown otherwise. If one sex strongly limits the reproductive capacity of the species or population, then only the decline in the limiting sex should be considered. We decided to restrict the automatic assignment of resilience categories in the Key Facts page to values of K , t_m and t_{max} and those records of fecundity estimates that referred to minimum number of eggs or pups per female per year, assuming that these were equivalent to average fecundity at first maturity (Musick 1999). Note that many small fishes may spawn several times per year (we exclude these for the time being) and large live bearers such as the coelacanth may have gestation periods of more than one year (we corrected fecundity estimates for those cases reported in the literature). Also, we excluded resilience estimates based on r_m (see below) as we are not yet confident with the reliability of the current method for estimating r_m . If users have independent r_m or fecundity estimates, they can refer to Table 1 for using this information."

Parameter	High	Medium	Low	Very low
Threshold	0.99	0.95	0.85	0.70
r_{max} (1/year)	> 0.5	0.16 – 0.50	0.05 – 0.15	< 0.05
K (1/year)	> 0.3	0.16 – 0.30	0.05 – 0.15	< 0.05
Fecundity (1/year)	> 10,000	100 – 1000	10 – 100	< 10
t_m (years)	< 1	2 – 4	5 – 10	> 10
t_{max} (years)	1 - 3	4 – 10	11 – 30	> 30

Taken from the FishBase manual, "Estimation of Life-History Key Facts":
<http://www.fishbase.us/manual/English/key%20facts.htm#resilience>