

# IFFO RS V2.0



## FISHERY ASSESSMENT METHODOLOGY AND TEMPLATE REPORT

Fishery Under Assessment	Chub Mackerel ( <i>Scomber Japonicus</i> )
Date	July 2017
Assessor	Deirdre Hoare

Application details and summary of the assessment outcome				
Name:				
Address:				
Country: Thailand, sourcing from Japan, China			Zip:	
Tel. No.		Fax. No.		
Email address:		Applicant Code		
Key Contact:		Title:		
Certification Body Details				
Name of Certification Body:		SAI Global (Ireland)		
Assessor Name	Peer Reviewer	Assessment Days	Initial/Surveillance/ Re-approval	Whole fish / By-product
Deirdre Hoare	Virginia Polonio	1	Surveillance	By-product
Assessment Period	2016			
Scope Details				
Management Authority (Country/State)		Sourced from Japan/China		
Main Species		Chub Mackerel ( <i>Scomber Japonicus</i> )		
Fishery Location		FAO 61 Pacific Northwest		
Gear Type(s)		Purse seine		
Outcome of Assessment				
Overall Outcome		Pass		
Clauses Failed		NA		
Peer Review Evaluation		Maintain approval		
Recommendation		Maintain approval		

**Assessment Determination**

This byproduct is imported to Thailand from Japan and China, FAO area 61. As stock assessment information is available for Chub mackerel in Japan it is assessed as category C, and although the information is slightly outdated, indications are of improvement in the SSB. Chub mackerel was assessed as category D for China as no stock assessment information could be found. The Productivity-Susceptibility Analysis found that this species is at low risk of overfishing. The IUCN Red List has categorised *Scomber japonicus* as 'Least Concern' in its global assessment.

**Peer Review Comments****Notes for On-site Auditor**

## Species-Specific Results

Category	Species	% landings	Outcome (Pass/Fail)	
Category A			A1	
			A2	
			A3	
			A4	
Category B				
Category C	Chub mackerel	NA	Pass	
Category D	Chub mackerel	NA	Pass	

[List all Category A and B species. List approximate total %age of landings which are Category C and D species; these do not need to be individually named here]

## HOW TO COMPLETE THIS ASSESSMENT REPORT

This assessment template uses a modular approach to assessing fisheries against the IFFO RS standard.

### By-products

The process for completing the template for **by-product raw material** is as follows:

1. ALL ASSESSMENTS: Complete the Species Characterisation table with the names of the by-product species and stocks under assessment. The '% landings' column can be left empty; all by-products are considered as Category C and D.
2. IF THERE ARE CATEGORY C BYPRODUCTS UNDER ASSESSMENT: Complete clause C1 for **each** Category C by-product.
3. IF THERE ARE CATEGORY D BYPRODUCTS UNDER ASSESSMENT: Complete Section D.
4. ALL OTHER SECTIONS CAN BE DELETED. Clauses M1 - M3, F1 - F3, and Sections A and B do not need to be completed for a by-product assessment.

By-product approval is awarded on a species-by-species basis. Each by-product species scoring a pass under the appropriate section may be approved against the IFFO RS Standard.

## SPECIES CATEGORISATION

The following table should be completed as fully as the available information permits. All species regularly\* caught in the fishery should be listed along with an estimate of the proportion of the catch each species represents. The species should then be divided into Type 1 and Type 2. **Type 1 species must represent 95% of the total catch. Type 2 species may represent a maximum of 5% of the catch (see Appendix B).**

\*Species which make up less than 0.1% of landings do not need to be listed (NOTE: ETP species are considered separately). The table should be extended if more space is needed. Discarded species should be included when known.

The 'stock' column should be used to differentiate when there are multiple biological or management stocks of one species captured by the fishery. The 'management' column should be used to indicate whether there is an adequate management regime specifically aimed at the individual species/stock. In some cases it will be immediately clear whether there is a species-specific management regime in place (for example, if there is an annual TAC). In less clear circumstances, the rule of thumb should be that if the species meets the minimum requirements of clauses A1-A4, an adequate species-specific management regime is in place.

NOTE: If any species is categorised as Endangered or Critically Endangered on the IUCN Red List, or if it appears in the CITES appendices, it **cannot** be approved for use as an IFFO RS raw material. This applied to whole fish as well as by-products.

### TYPE 1 SPECIES (Representing 95% of the catch or more)

**Category A:** Species-specific management regime in place.

**Category B:** No species-specific management regime in place.

### TYPE 2 SPECIES (Representing 5% OF THE CATCH OR LESS)

**Category C:** Species-specific management regime in place.

**Category D:** No species-specific management regime in place.

Common name	Latin name	Stock	% of landings	Management	Category
Chub mackerel	<i>Scomber japonicus</i>	Japan	NA	Yes	C
Chub mackerel	<i>Scomber japonicus</i>	China	NA	No	D

Category A species are assessed through an examination of the data collection, stock assessment, management measures, and stock status relating to the species. Category B species are assessed using a risk-based assessment covering similar areas. Category C species are assessed on stock status only. Category D species are assessed using a PSA analysis as described in the relevant section of this document.

## CATEGORY C SPECIES

In a whole fish assessment, Category C species are those which make up less than 5% of landings, but which are subject to a species-specific management regime. In most cases this will be because they are a commercial target in a fishery other than the one under assessment. In a by-product assessment, Category C species are those which are subject to a species-specific management regime, and are usually targeted species in fisheries for human consumption.

Clause C1 should be completed for **each** Category C species. If there are no Category C species in the fishery under assessment, this section can be deleted. A Category C species does not meet the minimum requirements of clause C1 should be re-assessed as a Category D species.

<b>Species Name</b>		Chub mackerel - Japan	
<b>C1</b>	<b>Category C Stock Status - Minimum Requirements</b>		
	C1.1	Fishery removals of the species in the fishery under assessment are included in the stock assessment process, OR are considered by scientific authorities to be negligible.	Yes
	C1.2	The species is considered, in its most recent stock assessment, to have a biomass above the limit reference point (or proxy), OR removals by the fishery under assessment are considered by scientific authorities to be negligible.	Yes
<b>Clause outcome:</b>			Pass
<b>Evidence</b> <p>Fishery removals of chub mackerel are included in the stock assessment process in Japan. Based on stock assessments of populations of this species in the Japan and the Tsushima Current between 1995 and 2005 (Watanabe 2009), SSB peaked in 1979 at 1,400,000 mt, and then declined to less than 38,000 mt in 2002, where it remained low but stable until 2004 when it increased to 300,000 mt in 2006 and then has slightly declined. In the Tsushima Current, SSB since 1973 averaged 350,000 mt, with a peak of 550,000 mt in 1989 and a low of 100,000 mt in 2004, and then has increased again to about 200,000 mt in 2006. Both stocks have increased in the past 10–12 years, likely due to better recruitment associated environmental changes and reduction in number of vessels and seasonal closures.</p> <p>In the most recent stock assessment available the SSB was found to be greater than <math>B_{lim}</math> which equals 450,000 tons (Fisheries Agency of Japan 2013). In 2012, SSB of this stock was 472,000 tons that was slightly higher than <math>B_{lim}</math> (Kawabata et al. 2014). The report states that the SSB should recover more, because current environmental conditions are good for this stock (Fisheries Agency of Japan, 2013).</p>			
<b>References</b> <p>Fisheries Research Agency Japan (2013) The population sizes of the Pacific stocks of Japanese sardine (<i>Sardinops melanostictus</i>) and chub mackerel (<i>Scomber japonicus</i>) are continuously increasing.  <a href="https://www.fra.affrc.go.jp/english/press/2013/20130625.html">https://www.fra.affrc.go.jp/english/press/2013/20130625.html</a></p>			
Standard clauses 1.3.2.1 - 1.3.2.4			

## CATEGORY D SPECIES

In a whole fish assessment, Category D species are those which make up less than 5% of landings and are not subject to a species-specific management regime. In the case of mixed trawl fisheries, Category D species may make up the majority of landings. In a by-product assessment, Category D species are those which are not subject to a species-specific management regime. In both cases, the comparative lack of scientific information on the status of the population of the species means that a risk-assessment style approach must be taken.

The process for assessing Category D species involves the use of a Productivity-Susceptibility Analysis (PSA) to further subdivide the species into 'Critical Risk', 'Major Risk' and 'Minor Risk' groups. If there are no Category D species in the fishery under assessment, this section can be deleted.

Productivity and susceptibility ratings are calculated using a process derived from the APFIC document "Regional Guidelines for the Management of Tropical Trawl Fisheries, which in turn was derived from papers by Patrick *et al* (2009) and Hobday *et al* (2007). Table D1 should be completed for each Category D species as follows:

- Firstly, the best available information should be used to fill in values for each productivity and susceptibility attribute.
- Table D2 should be used to convert each attribute value into a score between 1 and 3.
- The average score for productivity attributes and the average for susceptibility attributes should be calculated.
- Table D3 should be used to determine whether the species is required to meet the requirements of Table D4. A species which does not need to meet the requirements of D4 is automatically awarded a pass.
- Table D4 should be used to assess those species indicated by Table D3 to determine a pass/fail rating.
- Any Category D species which has been categorised by the IUCN Red List as Endangered or Critically Endangered, or which appears in the CITES appendices, automatically results in a fail.



<b>D1</b>	<b>Species Name</b>	<b>Chub mackerel China</b>	
	<b>Productivity Attribute</b>	<b>Value</b>	<b>Score</b>
	Average age at maturity (years)	2-4	2
	Average maximum age (years)	7	1
	Fecundity (eggs/spawning)	100,000 – 400,000	1
	Average maximum size (cm)	30	1
	Average size at maturity (cm)	26.1	1
	Reproductive strategy	Broadcast spawner	1
	Mean trophic level	3.4	3
	<b>Average Productivity Score</b>		<b>1.42</b>
	<b>Susceptibility Attribute</b>	<b>Value</b>	<b>Score</b>
	Overlap of adult species range with fishery	30	2
	Distribution	Global distribution	1
	Habitat	Pelagic-neritic	2
	Depth range	0-300	2
	Selectivity	1 to 2 times mesh size	2
	Post-capture mortality	Most dead, retained	3
	<b>Average Susceptibility Score</b>		<b>2</b>
	<b>PSA Risk Rating (From Table D3)</b>		<b>Pass</b>
	<b>Compliance rating</b>		<b>Pass</b>
	<b>References</b>		
	Fish Base <a href="http://fishbase.org/summary/Scomber-japonicus.html">http://fishbase.org/summary/Scomber-japonicus.html</a>		
	IUCN Redlist <a href="http://www.iucnredlist.org/details/170306/0">http://www.iucnredlist.org/details/170306/0</a>		
	<i>Standard clauses 1.3.2.1 - 1.3.2.4</i>		

Table D2 - Productivity / Susceptibility attributes and scores.

Productivity attributes	Low productivity/ High risk	Medium productivity/ Medium risk	High productivity/ Low risk
	Score 3	Score 2	Score 1
Average age at maturity (years)	>4	2 to 4	<2
Average maximum age (years)	>30	10 to 30	<10
Fecundity (eggs/spawning)	<1 000	1 000 to 10 000	>10 000
Average maximum size (cm)	>150	60 to 150	<60
Average size at maturity (cm)	>150	30 to 150	<30
Reproductive strategy	Live bearer, mouth brooder or significant parental investment	Demersal spawner "berried"	Broadcast spawner
Mean trophic level	>3.25	2.5–3.25	<2.5

Susceptibility attributes		High susceptibility/ High risk	Medium susceptibility/ Medium risk	Low susceptibility/ Low risk
		Score 3	Score 2	Score 1
Availability	1) Overlap of adult species range with fishery	>50% of stock occurs in the area fished	Between 25% and 50% of the stock occurs in the area fished	<25% of stock occurs in the area fished
	2) Distribution	Only in the country/ fishery	Limited range in the region	Throughout region/ global distribution
Encounterability	1) Habitat	Habitat preference of species make it highly likely to encounter trawl gear (e.g. demersal, muddy/sandy bottom)	Habitat preference of species make it moderately likely to encounter trawl gear (e.g. rocky bottom/reefs)	Depth or distribution of species make it unlikely to encounter trawl gear (e.g. epi-pelagic or meso-pelagic)
	2) Depth range	High overlap with trawl fishing gear (20 to 60 m depth)	Medium overlap with trawl fishing gear (10 to 20 m depth)	Low overlap with trawl fishing gear (0 to 10 m, >70 m depth)
Selectivity		Species >2 times mesh size or up to 4 m length	Species 1 to 2 times mesh size or 4 to 5 m length	Species <mesh size or >5 m length
Post capture mortality		Most dead or retained Trawl tow >3 hours	Alive after net hauled Trawl tow 0.5 to 3 hours	Released alive Trawl tow <0.5 hours

**Note:** Availability 2 is only used when there is no information for Availability 1; the most conservative score between Encounterability 1 and 2 is used.

<b>D3</b>		<b>Average Susceptibility Score</b>		
		<b>1 - 1.75</b>	<b>1.76 - 2.24</b>	<b>2.25 - 3</b>
<b>Average Productivity Score</b>	<b>1 - 1.75</b>	PASS	PASS	PASS
	<b>1.76 - 2.24</b>	PASS	PASS	TABLE D4
	<b>2.25 - 3</b>	PASS	TABLE D4	TABLE D4

## **SOCIAL CRITERION**

In addition to the scored criteria listed above, applicants must commit to ensuring that vessels operating in the fishery adhere to internationally recognised guidance on human rights. They must also commit to ensuring there is no use of enforced or unpaid labour in the fleet(s) operating upon the resource.

## Appendix A - Determining Resilience Ratings

The assessment of Category B species described in this assessment report template utilises a resilience rating system suggested by the American Fisheries Society. This approach was chosen because it is also used by FishBase, and so the resilience ratings for many thousands of species are freely available online. As described by FishBase, the following is the process used to arrive at the resilience ratings:

*“The American Fisheries Society (AFS) has suggested values for several biological parameters that allow classification of a fish population or species into categories of high, medium, low and very low resilience or productivity (Musick 1999). If no reliable estimate of  $r_m$  (see below) is available, the assignment is to the lowest category for which any of the available parameters fits. For each of these categories, AFS has suggested thresholds for decline over the longer of 10 years or three generations. If an observed decline measured in biomass or numbers of mature individuals exceeds the indicated threshold value, the population or species is considered vulnerable to extinction unless explicitly shown otherwise. If one sex strongly limits the reproductive capacity of the species or population, then only the decline in the limiting sex should be considered. We decided to restrict the automatic assignment of resilience categories in the Key Facts page to values of  $K$ ,  $t_m$  and  $t_{max}$  and those records of fecundity estimates that referred to minimum number of eggs or pups per female per year, assuming that these were equivalent to average fecundity at first maturity (Musick 1999). Note that many small fishes may spawn several times per year (we exclude these for the time being) and large live bearers such as the coelacanth may have gestation periods of more than one year (we corrected fecundity estimates for those cases reported in the literature). Also, we excluded resilience estimates based on  $r_m$  (see below) as we are not yet confident with the reliability of the current method for estimating  $r_m$ . If users have independent  $r_m$  or fecundity estimates, they can refer to Table 1 for using this information.”*

Parameter	High	Medium	Low	Very low
Threshold	0.99	0.95	0.85	0.70
$r_{max}$ (1/year)	> 0.5	0.16 - 0.50	0.05 - 0.15	< 0.05
$K$ (1/year)	> 0.3	0.16 - 0.30	0.05 - 0.15	< 0.05
Fecundity (1/year)	> 10,000	100 - 1000	10 - 100	< 10
$t_m$ (years)	< 1	2 - 4	5 - 10	> 10
$t_{max}$ (years)	1 - 3	4 - 10	11 - 30	> 30

[Taken from the FishBase manual, “Estimation of Life-History Key Facts”,  
<http://www.fishbase.us/manual/English/key%20facts.htm#resilience>]

## Appendix B – Background on the 5% catch rule

The proposed fishery assessment methodology uses a species categorisation approach to divide the catch in the assessment fishery into groups. These groups are:

- **Category A:** “Target” species with a species-specific management regime in place.
- **Category B:** “Target” species with no species-specific management regime in place.
- **Category C:** “Non-target” species with a species-specific management regime in place.
- **Category D:** “Non-target” species with no species-specific management regime in place

The distinction between 'target' and 'non-target' species is made to enable the assessment to consider the impact of the fishery on all the species caught regularly, without requiring a full assessment be conducted for each. Thus 'target' species are subjected to a more detailed assessment, while 'non-target' species are considered more briefly. For the purposes of the IFFO RS fishery assessment, 'target' and 'non-target' species are defined by their prevalence in the catch, by weight. Applicants must declare which species are considered 'target' species in the fishery, and the combined weight of these must be at least 95% of the annual catch. The remaining 5% can be made up of 'non-target' species. Note also that ETP species are considered separately, irrespective of their frequency of occurrence in the catch.

The proposed use of 5% as a limit for 'non-target' species is one area in which feedback is being sought via the public consultation. The decision to propose a value of 5% ensures consistency with other fishery assessment programmes, such as the MSC which uses 5% to distinguish between 'main' and 'minor' species (see MSC Standard, SA3.4 and GSA3.4.2); and Seafood Watch, which uses 5% when defining the 'main' species for the assessment (see Seafood Watch Standard, Criterion 2). The value is also consistent with the approach used in Version 1 of the IFFO RS Standard, in which up to 5% of the raw material could be comprised of 'unassessed' species.

**Comments on this proposition are welcomed along with any other feedback on the proposed approach.**