

MarinTrust Standard V2

By-product Fishery Assessment John Dory (Zeus faber) FAO Area 27 North East Atlantic excluding Division 7.a

MarinTrust Programme

Unit C, Printworks 22 Amelia Street London

SE17 3BZ

E: standards@marin-trust.com

T: +44 2039 780 819



Table 1 Application details and summary of the assessment outcome

	Species:	John Dory (Zeus faber)		
	Geographical area:	FAO Area 27 North East Atlantic		
Fishery Under Assessment	Country of origin of the product:	Flag country not supplied by client		
	Stock:	John Dory in FAO 27 excluding Division 7.a		
Date	19 July 2022			
Report Code	FRA09			
Assessor	Matthew Jew			
Country of origin of the product - PASS	Flag country not supplied by client			
Country of origin of the product - FAIL	NA			

Application details and summary of the assessment outcome							
Company Name(s): Bioceval							
Country: France							
Email address:	Email address: Applicant Code:						
Certification Body Det	ails						
Name of Certification	Body:	Global Trust C	Global Trust Certification				
Assessor	Peer Reviewer	Assessment Days	Initial/Surveillance/ Re-approval				
Matthew Jew	Léa Lebechnech	1	Re-Approval				
Assessment Period	Up to July 2022						

Scope Details	
Main Species	John Dory (Zeus faber)
Stock	FAO Area 27 Northeast Atlantic
Fishery Location	John Dory in FAO Area 27 excluding Division 7.a
Management Authority	European Union (Common Fisheries Policy)
(Country/ State)	Laropean official (common risheries rolley)
Gear Type(s)	Demersal trawls, seines, gillnets, and beam trawls
Outcome of Assessment	
Peer Review Evaluation	Agree with the assessor's determination
Recommendation	APPROVED

Table 2. Assessment Determination

Assessment Determination

If any species is categorised as Endangered or Critically Endangered on IUCN's Red List, or if it appears in the CITES appendices, it cannot be approved for use as Marin trust raw material. John Dory (*Zeus faber*) do not appear as Endangered or Critically Endangered on IUCN's Red List, nor do they appear in CITES appendices; therefore, *Zeus faber* is eligible for approval for use as Marin trust by-product raw material.

One stock is part of this assessment:

1) Zeus faber in FAO 27 excluding 7a North-eastern Atlantic Ocean.

There are no management measures specifically implemented for the John Dory stock. The species is mainly caught in mixed trawl fisheries. There are no species-specific reference points. Therefore, following MarinTrust criteria, the species is assessed as Category D.

Table D1 (PSA) shows that the stock as an average productivity score of 1.71 and an average susceptibility score of 2.75. The PSA risk rating results (Table D3) determined that the species passes.

Therefore, John Dory in FAO 27 excluding 7a is **APPROVED** for the production of fishmeal and fish oil under the current MarinTrust v2.0 by-products standard.

Fishery Assessment Peer Review Comments

The assessor correctly classified John Dory in the FAO Area 27 (Northeast Atlantic excl. Division 7.a) stock as category D, as reference points are undefined to assess status of the stock relative to.

With an average productivity score of 1.71 and an average susceptibility score of 2.75, it passes as per Table D3.

Therefore, John Dory in the FAO Area 27 (Northeast Atlantic excl. Division 7.a) is APPROVED.

Notes for On-site Auditor

Determine which flag state(s) the plant is sourcing its John Dory from.



Species Categorisation

NB: If any species is categorised as Endangered or Critically Endangered on the IUCN Red List, or if it appears in CITES Appendix 1, it **cannot** be approved for use as an MarinTrust raw material.

IUCN Red list Category

By-product material from a species listed by IUCN (the International Union for Conservation of Nature) under the Red List for the following categories shall immediately fail the assessment;

- EXTINCT (E) AND EXTINCT IN THE WILD (EW)
- CRITICALLY ENDANGERED (CR) facing an extremely high risk of extinction in the wild.
- ENDANGERED (EN) facing a very high risk of extinction in the wild.

By-product material may be used from the following categories provided that all clauses in the MarinTrust standard are passed.

- VULNERABLE (VU) facing a high risk of extinction in the wild.
- NEAR THREATENED (NT) does not qualify for above now, but is close or is likely to qualify for, a threatened category in the near future.
- LEAST CONCERN (LC) Widespread and abundant.
- DATA DEFICIENT (DD) and NOT EVALUATED (NE)

Table 3 Species Categorisation Table

Common name	Latin name	Stock	Management	Category	IUCN Red List Category ¹	CITES Appendix 1 ²
John Dory	Zeus faber	John Dory in FAO 27 excluding Division 7.a	EU (Common Fisheries Policy)	D	DD	No

¹ https://www.iucnredlist.org/

² https://cites.org/eng/app/appendices.php

CATEGORY D SPECIES

Category D species are those which are not subject to a species-specific management regime. In the case of mixed trawl fisheries, Category D species may make up the majority of landings. The comparative lack of scientific information on the status of the population of the species means that a risk-assessment style approach must be taken.

Species Name		John Dory (Zeus faber)				
Productivity Attı	ribute	Value	Score			
Average age at maturity (years)		4	1			
Average maximum age (years)		12	2			
Fecundity (eggs/spawning)		See note below	3			
Average maximum size (cm)		90	1			
Average size at maturity (cm)		35.1	1			
Reproductive strategy		Pelagic Spawn (Broadcast spawner)	1			
Mean trophic level		4.5	3			
		Average Productivity Score	1.71			
Susceptibility Att	ribute	Value	Score			
Availability (area overlap)		25%	2			
Encounterability (the position of within the water column relative		High overlap with fishing gear	3			
Selectivity of gear type		See note below	3			
Post-capture mortality		Retained Species	3			
		Average Susceptibility Score	2.75			
		PSA Risk Rating (From Table D3)	Pass			
		Compliance rating	PASS			

Further justification for productivity scoring (where relevant)

There is no published information regarding the fecundity of this species or its close relatives. Fecundity was given maximum score as a conservative precaution.

Further justification for susceptibility scoring (where relevant)

According to the map below, the area overlap of the fishery with the species can be estimated as around 25%.



This map was computer-generated and has not yet been reviewed.

Zeus faber AquaMaps Data sources: GBIF OBIS

This species resides in the demersal environment (close to the sea bed) and is often encountered with the bottom gear used in the fishery, which is why it was given the value of high overlap for encounterability. There is limited information about the selectivity of gear type as it pertains to the average size at maturity for the species. 'Selectivity of gear type' was scored a 3 as a conservative precaution.

References



FAO. 2022. Zeus faber Linnaeus, 1758. Fisheries and Aquaculture Division [online]. Rome. https://www.fao.org/fishery/en/aqspecies/2250/en. Access on 20 July 2022.

Froese, R. and D. Pauly. Editors. 2022. FishBase. Life History of *Zeus faber* (John Dory). https://www.fishbase.de/summary/zeus-faber.html. Accessed 20 July 2022.

Iwamoto, T. 2015. *Zeus faber*. The IUCN Red List of Threatened Species 2015: e.T198769A42390771. https://dx.doi.org/10.2305/IUCN.UK.2015-4.RLTS.T198769A42390771.en. Accessed on 20 July 2022.

Standard clauses 1.3.2.2



Table D2 - Productivity / Susceptibility attributes and scores.

Productivity attributes	High productivity (Low risk, score = 1)	Medium productivity (medium risk, score = 2)	Low productivity (high risk, score = 3)
Average age at maturity	<5 years	5-15 years	>15 years
Average maximum age	<10 years	10-25 years	>25 years
Fecundity	>20,000 eggs per year	100-20,000 eggs per year	<100 eggs per year
Average maximum size	<100 cm	100-300 cm	>300 cm
Average size at maturity	<40 cm	40-200 cm	>200 cm
Reproductive strategy	Broadcast spawner	Demersal egg layer	Live bearer
Mean Trophic Level	<2.75	2.75-3.25	>3.25

Susceptibility attributes		ow susceptibility ow risk, score = 1)	Medium susceptibility (medium risk, score = 2)		High susceptibility (high risk, score = 3)		
Areal overlap (availability) Overlap of the fishing effort with the species range	<10% overlap		10	10-30% overlap		>30% overlap	
Encounterability The position of the stock/species within the water column relative to the fishing gear, and the position of the stock/species within the habitat relative to the position of the gear	fis	ow overlap with hing gear (low acounterability).	Medium overlap with fishing gear.		High overlap with fishing gear (high encounterability). Default score for target species		
Selectivity of gear type Potential of the gear to retain species	а	Individuals < size at maturity are rarely caught	a	Individuals < size at maturity are regularly caught.	a	Individuals < size at maturity are frequently caught	
	b	Individuals < size at maturity can escape or avoid gear.	b	Individuals < half the size at maturity can escape or avoid gear.	b	Individuals < half the size at maturity are retained by gear.	
Post-capture mortality (PCM) The chance that, if captured, a species would be released and that it would be in a condition permitting subsequent survival	Evidence of majority released post-capture and survival.		Evidence of some released post-capture and survival.		Retained species or majority dead when released.		



D3		Average Susceptibility Score				
		1 - 1.75	1.76 - 2.24	4 2.25 - 3		
Average Productivity	1 - 1.75	PASS	PASS	PASS		
Score	1.76 - 2.24	PASS	PASS	TABLE D4		
	2.25 - 3	PASS	TABLE D4	TABLE D4		