



RESPONSIBLE
SUPPLY

IFFO RS
Global Standard for Responsible Supply
of Marine Ingredients

IFFO RS Limited

T: +44 (0) 2030 539 195
E: Standards@iffors.com
W: www.iffors.com

Unit C, Printworks | 22 Amelia Street
London, SE17 3BZ | United Kingdom



RESPONSIBLE
SUPPLY

IFFO
RS

ASSURED



Global Standard for Responsible Supply of Marine Ingredients Fishery Assessment Methodology and Template Report V2.0



RESPONSIBLE
SUPPLY

IFFO RS
Global Standard for Responsible Supply
of Marine Ingredients



Fishery Under Assessment	Brazilian Flathead, Pez palo (<i>Percophis Brasiliensis</i>)
Date	February 2019
Assessor	V.Polonio

Application details and summary of the assessment outcome				
Name: Coomarpes				
Address:				
Country: Argentina		Zip:		
Tel. No.:		Fax. No.:		
Email address:		Applicant Code		
Key Contact:		Title:		
Certification Body Details				
Name of Certification Body:		SAI Global Ltd		
Assessor Name	Peer Reviewer	Assessment Days	Initial/Surveillance/Re-approval	Whole fish/ By-product
V. Polonio	J. Daly	0.5	Surveillance 1	By-product
Assessment Period		2018		

Scope Details	
Management Authority (Country/State)	Argentina
Main Species	Brazilian Flathead (<i>Percophis Brasiliensis</i>)
Fishery Location	Atlantic Southwest FAO Zone NR 41°S
Gear Type(s)	Trawl
Outcome of Assessment	
Overall Outcome	Pass
Clauses Failed	None
Peer Review Evaluation	AGREE
Recommendation	APPROVE

Assessment Determination
<p>There is no new stock assessment since 2016 and therefore information from the last assessment has not shown relevant changes. However there had been a workshop between INIPED and SIMPLOT to develop a new model that can correct uncertainties from the last stock assessment.</p> <p>This species has not yet been assessed for the IUCN Red List (website accessed 09.02.19). Brazilian Flathead (<i>Percophis Brasiliensis</i>) is approved by the assessment team for the production of fishmeal and fish oil under the IFFO-RS v 2.0 standard (by-products).</p>
Peer Review Comments
Agree
Notes for On-site Auditor

Note: This table should be completed for whole fish assessments only.

Species-Specific Results

Category	Species	% landings	Outcome (Pass/Fail)	
Category A			A1	
			A2	
			A3	
			A4	
Category B				
Category C	Brazilian Flathead, Pez palo (<i>Percophis Brasiliensis</i>)	N/A	PASS	
Category D				

[List all Category A and B species. List approximate total %age of landings which are Category C and D species; these do not need to be individually named here]

HOW TO COMPLETE THIS ASSESSMENT REPORT

This assessment template uses a modular approach to assessing fisheries against the IFFO RS standard.

Whole Fish

The process for completing the template for a **whole fish** assessment is as follows:

1. ALL ASSESSMENTS: Complete the Species Characterisation table, to determine which categories of species are present in the fishery.
2. ALL ASSESSMENTS: Complete clauses M1, M2, M3: Management.
3. IF THERE ARE CATEGORY A SPECIES IN THE FISHERY: Complete clauses A1, A2, A3, A4 for **each** Category A species.
4. IF THERE ARE CATEGORY B SPECIES IN THE FISHERY: Complete the Section B risk assessment for **each** Category B species.
5. IF THERE ARE CATEGORY C SPECIES IN THE FISHERY: Complete clause C1 for **each** Category C species.
6. IF THERE ARE CATEGORY D SPECIES IN THE FISHERY: Complete Section D.
7. ALL ASSESSMENTS: Complete clauses F1, F2, F3: Further Impacts.

A fishery must score a pass in **all applicable clauses** before approval may be recommended. To achieve a pass in a clause, the fishery/species must meet **all** of the minimum requirements.

By-products

The process for completing the template for **by-product raw material** is as follows:

1. ALL ASSESSMENTS: Complete the Species Characterisation table with the names of the by-product species and stocks under assessment. The ‘% landings’ column can be left empty; all by-products are considered as Category C and D.
2. IF THERE ARE CATEGORY C BYPRODUCTS UNDER ASSESSMENT: Complete clause C1 for **each** Category C by-product.
3. IF THERE ARE CATEGORY D BYPRODUCTS UNDER ASSESSMENT: Complete Section D.
4. ALL OTHER SECTIONS CAN BE DELETED. Clauses M1 - M3, F1 - F3, and Sections A and B do not need to be completed for a by-product assessment.

By-product approval is awarded on a species-by-species basis. Each by-product species scoring a pass under the appropriate section may be approved against the IFFO RS Standard.

SPECIES CATEGORISATION

The following table should be completed as fully as the available information permits. Any species representing more than 0.1% of the annual catch should be listed, along with an estimate of the proportion of the catch each species represents. The species should then be divided into Type 1 and Type 2 as follows:

- **Type 1 Species** can be considered the ‘target’ or ‘main’ species in the fishery. They make up the bulk of annual landings and are subjected to a detailed assessment.
- **Type 2 Species** can be considered the ‘bycatch’ or ‘minor’ species in the fishery. They make up a small proportion of the annual landings and are subjected to relatively high-level assessment.

Type 1 Species must represent 95% of the total annual catch. Type 2 Species may represent a maximum of 5% of the annual catch (see Appendix B).

Species which make up less than 0.1% of landings do not need to be listed (NOTE: ETP species are considered separately). The table should be extended if more space is needed. Discarded species should be included when known.

The ‘stock’ column should be used to differentiate when there are multiple biological or management stocks of one species captured by the fishery. The ‘management’ column should be used to indicate whether there is an adequate management regime specifically aimed at the individual species/stock. In some cases it will be immediately clear whether there is a species-specific management regime in place (for example, if there is an annual TAC). In less clear circumstances, the rule of thumb should be that if the species meets the minimum requirements of clauses A1-A4, an adequate species-specific management regime is in place.

NOTE: If any species is categorised as Endangered or Critically Endangered on the IUCN Red List, or if it appears in the CITES appendices, it **cannot** be approved for use as an IFFO RS raw material. This applied to whole fish as well as by-products.

TYPE 1 SPECIES (Representing 95% of the catch or more)

Category A: Species-specific management regime in place.

Category B: No species-specific management regime in place.

TYPE 2 SPECIES (Representing 5% OF THE CATCH OR LESS)

Category C: Species-specific management regime in place.

Category D: No species-specific management regime in place.

Common name	Latin name	Stock	% of landings	Management	Category
Brazilian Flathead, Pez palo	<i>Percophis Brasiliensis</i>	Mar del Plata	N/A	Argentina	C

CATEGORY C SPECIES

In a whole fish assessment, Category C species are those which make up less than 5% of landings, but which are subject to a species-specific management regime. In most cases this will be because they are a commercial target in a fishery other than the one under assessment. In a by-product assessment, Category C species are those which are subject to a species-specific management regime, and are usually targeted species in fisheries for human consumption.

Clause C1 should be completed for **each** Category C species. If there are no Category C species in the fishery under assessment, this section can be deleted. A Category C species does not meet the minimum requirements of clause C1 should be re-assessed as a Category D species.

Species Name		Brazilian flathead <i>Percophis Brasiliensis</i>																										
C1	Category C Stock Status - Minimum Requirements																											
	C1.1	Fishery removals of the species in the fishery under assessment are included in the stock assessment process, OR are considered by scientific authorities to be negligible.	Yes																									
	C1.2	The species is considered, in its most recent stock assessment, to have a biomass above the limit reference point (or proxy), OR removals by the fishery under assessment are considered by scientific authorities to be negligible.	Yes																									
Clause outcome:				Pass																								
Evidence																												
<p>C1.1 Fishery removals of the species in the fishery under assessment are included in the stock assessment process, OR are considered by scientific authorities to be negligible.</p> <p>Fishery removals of the species in the Argentine Brazilian flathead fishery are included in the stock assessment process. The stock assessment is conducted by the National Institute for Fisheries Research and Development (Instituto Nacional de Investigacion y Desarrollo Pesquero, or INIDEP) and considers the fishery operating North of 41°S Argentina. Further data from a workshop carried out in 2018 from Uruguay has been considered. Illegal catches are estimated in the models from 2009.</p> <p>Fishery removals of the species in the fishery under assessment are included in the stock assessment process.</p> <p>The species passes Clause C1.1</p>																												
<p>C1.2 The species is considered, in its most recent stock assessment, to have a biomass above the limit reference point (or proxy), OR removals by the fishery under assessment are considered by scientific authorities to be negligible.</p> <p>The last stock assessment was done in 2016. The results have shown that Spawning stock biomass is above B_{lim} and B_{msy}. The stock assessment uses well-accepted methods, including several surveys of different population segments that provide fishery-independent data. However, estimates are highly variable due to the uncertainty in the model parameters. IUU fishing, foreign vessels catches, discards and bycatch estimates are included in the assessment since 2009. Different models were used to estimate stock status, all of them have shown uncertainties and different estimation. The main results are shown in Table 1. The species passes Clause C1.2.</p>																												
<p>Table 1. Results of the estimation of the acceptable biological catch (ABC) for the different evaluation models. R2</p>																												
<table border="1"> <thead> <tr> <th rowspan="2">Modelo</th> <th colspan="2" rowspan="2">Aplicación</th> <th rowspan="2">Punto de Referencia</th> <th colspan="2">Argentina</th> </tr> <tr> <th>CPUE</th> <th>CPUE+Campaña</th> </tr> </thead> <tbody> <tr> <td rowspan="2">Modelos no estructurados</td> <td rowspan="2">Schaefer</td> <td>Máxima Verosimilitud</td> <td>RMS</td> <td>6.799</td> <td>6.613</td> </tr> <tr> <td>Inferencia Bayesiana</td> <td>RMS</td> <td>5.316</td> <td>-</td> </tr> <tr> <td>Modelo estructurado</td> <td>ASPM</td> <td>Máxima Verosimilitud</td> <td>$BRLP > 0,3 BRV$</td> <td>8.097</td> <td>3.623</td> </tr> </tbody> </table>					Modelo	Aplicación		Punto de Referencia	Argentina		CPUE	CPUE+Campaña	Modelos no estructurados	Schaefer	Máxima Verosimilitud	RMS	6.799	6.613	Inferencia Bayesiana	RMS	5.316	-	Modelo estructurado	ASPM	Máxima Verosimilitud	$BRLP > 0,3 BRV$	8.097	3.623
Modelo	Aplicación		Punto de Referencia	Argentina																								
				CPUE	CPUE+Campaña																							
Modelos no estructurados	Schaefer	Máxima Verosimilitud	RMS	6.799	6.613																							
		Inferencia Bayesiana	RMS	5.316	-																							
Modelo estructurado	ASPM	Máxima Verosimilitud	$BRLP > 0,3 BRV$	8.097	3.623																							
<p>References:</p> <p>R1 Fishbase Brazilian flathead: http://www.fishbase.org/summary/Percophis-brasiliensis.html</p> <p>R2 Instituto Nacional de Investigación y Desarrollo Pesquero (INIDEP) https://www.argentina.gob.ar/inidep</p>																												

Appendix A - Determining Resilience Ratings

The assessment of Category B species described in this assessment report template utilises a resilience rating system suggested by the American Fisheries Society. This approach was chosen because it is also used by FishBase, and so the resilience ratings for many thousands of species are freely available online. As described by FishBase, the following is the process used to arrive at the resilience ratings:

“The American Fisheries Society (AFS) has suggested values for several biological parameters that allow classification of a fish population or species into categories of high, medium, low and very low resilience or productivity (Musick 1999). If no reliable estimate of r_m (see below) is available, the assignment is to the lowest category for which any of the available parameters fits. For each of these categories, AFS has suggested thresholds for decline over the longer of 10 years or three generations. If an observed decline measured in biomass or numbers of mature individuals exceeds the indicated threshold value, the population or species is considered vulnerable to extinction unless explicitly shown otherwise. If one sex strongly limits the reproductive capacity of the species or population, then only the decline in the limiting sex should be considered. We decided to restrict the automatic assignment of resilience categories in the Key Facts page to values of K , t_m and t_{max} and those records of fecundity estimates that referred to minimum number of eggs or pups per female per year, assuming that these were equivalent to average fecundity at first maturity (Musick 1999). Note that many small fishes may spawn several times per year (we exclude these for the time being) and large live bearers such as the coelacanth may have gestation periods of more than one year (we corrected fecundity estimates for those cases reported in the literature). Also, we excluded resilience estimates based on r_m (see below) as we are not yet confident with the reliability of the current method for estimating r_m . If users have independent r_m or fecundity estimates, they can refer to Table 1 for using this information.”

Parameter	High	Medium	Low	Very low
Threshold	0.99	0.95	0.85	0.70
r_{max} (1/year)	> 0.5	0.16 – 0.50	0.05 – 0.15	< 0.05
K (1/year)	> 0.3	0.16 – 0.30	0.05 – 0.15	< 0.05
Fecundity (1/year)	> 10,000	100 – 1000	10 – 100	< 10
t_m (years)	< 1	2 – 4	5 – 10	> 10
t_{max} (years)	1 - 3	4 – 10	11 – 30	> 30

Taken from the FishBase manual, “Estimation of Life-History Key Facts”:

<http://www.fishbase.us/manual/English/key%20facts.htm#resilience>]

Appendix B – Background on the 5% catch rule

The proposed fishery assessment methodology uses a species categorisation approach to divide the catch in the assessment fishery into groups. These groups are:

- **Category A:** “Target” species with a species-specific management regime in place.
- **Category B:** “Target” species with no species-specific management regime in place.
- **Category C:** “Non-target” species with a species-specific management regime in place.

- **Category D:** “Non-target” species with no species-specific management regime in place

The distinction between 'target' and 'non-target' species is made to enable the assessment to consider the impact of the fishery on all the species caught regularly, without requiring a full assessment be conducted for each. Thus 'target' species are subjected to a more detailed assessment, while 'non-target' species are considered more briefly. For the purposes of the IFFO RS fishery assessment, 'target' and 'non-target' species are defined by their prevalence in the catch, by weight. Applicants must declare which species are considered 'target' species in the fishery, and the combined weight of these must be at least 95% of the annual catch. The remaining 5% can be made up of 'non-target' species. Note also that ETP species are considered separately, irrespective of their frequency of occurrence in the catch.

The proposed use of 5% as a limit for 'non-target' species is one area in which feedback is being sought via the public consultation. The decision to propose a value of 5% ensures consistency with other fishery assessment programmes, such as the MSC which uses 5% to distinguish between 'main' and 'minor' species (see MSC Standard, SA3.4 and GSA3.4.2); and Seafood Watch, which uses 5% when defining the 'main' species for the assessment (see Seafood Watch Standard, Criterion 2). The value is also consistent with the approach used in Version 1 of the IFFO RS Standard, in which up to 5% of the raw material could be comprised of 'unassessed' species.

Comments on this proposition are welcomed along with any other feedback on the proposed approach.