

## MarinTrust Standard V2

## By-product Fishery Assessment Report Template

#### **MarinTrust Programme**

Unit C, Printworks 22 Amelia Street London SE17 3BZ

E: standards@marin-trust.com

T: +44 2039 780 819



# Table 1 Application details and summary of the assessment outcome

	Species:	Mackerel (Scomber scombrus)	
	Geographical area:	FAO Area 27 North East Atlantic	
Fishery Under Assessment	Country of origin of the product:	Iceland	
	Stock:	Mackerel in ICES Division 5.a	
Date	18 January 2022		
Report Code	BP 009		
Assessor	Geraldine Criquet		
Country of origin of the product - PASS	Iceland		
Country of origin of the product - FAIL	NA		

Application details and summary of the assessment outcome						
Name: Síldarvinnslan hf, Vinnslustöðin hf.						
Address:						
Country: Iceland		Zip:				
Tel. No.:		Fax. No.:				
Email address:		Applicant Cod	e:			
Key Contact:		Title:				
<b>Certification Body Det</b>	ails					
Name of Certification	Body:	Global Trust Certification				
Assessor Peer Reviewer		Assessment Days	Initial/Surveillance/ Re-approval			
Geraldine Criquet	Conor Donnelly	0.5	Surveillance 1			
Assessment Period To January 2022						

Scope Details		
Main Species Mackerel (Scomber scombrus)		
Stock	Mackerel in ICES Division 5.a	
Fishery Location	FAO Area 27 Northeast Atlantic Ocean	
Management Authority (Country/ State)	Iceland management authority	
Gear Type(s)	Diverse fishing gears	
Peer Review Evaluation	Agree with recommendation	
Recommendation	APPROVED	



## Table 2. Assessment Determination

#### **Assessment Determination**

If any species is categorised as Endangered or Critically Endangered on IUCN's Red List, or if it appears in the CITES appendices, it cannot be approved for use as Marin Trust raw material. Mackerel (*Scomber scombrus*) is neither listed as Endangered or Critically Endangered on IUCN's Red List, nor listed in CITES appendices; therefore, mackerel is eligible for approval for use as Marin Trust by-product raw material.

There is no long-term management strategy for this stock. Reference points are defined for the stock, therefore it was assessed under category C.

Fishery removals are included in the stock assessment process, it PASSES Clause C1.1. The stock is considered, in its most recent stock assessment, to have a biomass above the limit reference point, it PASSES Clause C1.2.

Therefore, mackerel in ICES Division 5.a is **APPROVED** for the production of fishmeal and fish oil under the current Marin Trust v 2.0 by-products.

#### **Fishery Assessment Peer Review Comments**

The stock has been correctly defined as a Category C stock. It also passes clause C1.1 and C1.2 as catches are taken into account in the model and the latest stock biomass is above target and limit reference points. Consequently, mackerel in ICES Division 5.a should be approved for the production of fishmeal and fish oil under the current Marin Trust v 2.0 by-products

Notes for On-site Auditor



## **Species Categorisation**

**NB:** If any species is categorised as Endangered or Critically Endangered on the IUCN Red List, or if it appears in CITES Appendix 1, it **cannot** be approved for use as an MARINTRUST raw material.

#### **IUCN Redlist Category**

By-product material from a species listed by IUCN (the International Union for Conservation of Nature) under the Red List for the following categories shall immediately fail the assessment;

- EXTINCT (E) AND EXTINCT IN THE WILD (EW)
- CRITICALLY ENDANGERED (CR) facing an extremely high risk of extinction in the wild.
- ENDANGERED (EN) facing a very high risk of extinction in the wild.

By-product material may be used from the following categories provided that all clauses in the MarinTrust standard are passed.

- VULNERABLE (VU) facing a high risk of extinction in the wild.
- NEAR THREATENED (NT) does not qualify for above now, but is close or is likely to qualify for, a threatened category in the near future.
- LEAST CONCERN (LC) Widespread and abundant.
- DATA DEFICIENT (DD) and NOT EVALUATED (NE)

## Table 3 Species Categorisation Table

Common name	Latin name	Stock	Management	Category	IUCN Red List Category <sup>1</sup>	CITES Appendix 1 <sup>2</sup>
Mackerel	Scomber scombrus	Mackerel in ICES Division 5.a	Iceland	С	LC	No

<sup>&</sup>lt;sup>1</sup> <u>https://www.iucnredlist.org/</u>

<sup>&</sup>lt;sup>2</sup> https://cites.org/eng/app/appendices.php

### **CATEGORY C SPECIES**

In a whole fish assessment, Category C species are those which make up less than 5% of landings, but which are subject to a species-specific management regime. In most cases this will be because they are a commercial target in a fishery other than the one under assessment.


Clause C1 should be completed for **each** Category C species. If there are no Category C species in the fishery under assessment, this section can be deleted. Where a species fails this Clause, it should be assessed as a Category D species instead.

Spe	ecies	Name	Mackerel (Scomber scombrus)			
<b>C1</b>	Catego	ory C Stock Sta	atus - Minimum Requirements			
CI	C1.1	Fishery remo	ovals of the species in the fishery under assessment are included in the stock	Yes		
		assessment	process, OR are considered by scientific authorities to be negligible.			
	C1.2 The species is considered, in its most recent stock assessment, to have a biomass above the					
	limit reference point (or proxy), OR removals by the fishery under assessment are					
	<u> </u>		Clause outcome:	PASS		

C1.1 Fishery removals of the species in the fishery under assessment are included in the stock assessment process, OR are considered by scientific authorities to be negligible.

The stock assessment type is an age-based analytical model (SAM) that uses catches in the model and the forecast. Catches are presented in Figure 1.

Therefore, fishery removals of the stock, including from the fishery under assessment, are included in the stock assessment process, it PASSES Clause C1.1



**Figure 1**. Mackerel in Division 5.a. Long-term trends in catches; the yellow shaded catches prior to 2000 are the years that have been down-weighted in the assessment because of the considerable underreporting that is suspected to have taken place.

C1.2 The species is considered, in its most recent stock assessment, to have a biomass above the limit reference point (or proxy), OR removals by the fishery under assessment are considered by scientific authorities to be negligible.

The spawning-stock size is above MSY Btrigger, Bpa and Blim (Figure 2).



Therefore, the stock is considered, in its most recent stock assessment, to have a biomass above the limit reference point, **C1.2** is met.




Figure 2. Mackerel in ICES Division 5.a. Spawning stock biomass.

#### References

Iceland Marine and freshwater Research Institute 2021. Status of Marine Stocks and Advice 2021. Mackerel, 30 September 2021. <a href="https://www.hafogvatn.is/static/extras/images/makrill">https://www.hafogvatn.is/static/extras/images/makrill</a> 20211278360.pdf

Collette, B., Fernandes, P. & Heessen, H. 2015. Scomber scombrus. The IUCN Red List of Threatened Species 2015: e.T170354A18207463. Accessed on 18 January 2022.

https://www.iucnredlist.org/species/170354/18207463

Links	
MARINTRUST Standard clause	1.3.2.2
FAO CCRF	7.5.3
GSSI	D.3.04, D5.01



## **CATEGORY D SPECIES**

Category D species are those which make up less than 5% of landings and are not subject to a species-specific management regime. In the case of mixed trawl fisheries, Category D species may make up the majority of landings. The comparative lack of scientific information on the status of the population of the species means that a risk-assessment style approach must be taken.

D1	<b>Species Name</b>				
	Productivity Attribut	te Value	Score		
	Average age at maturity (years)				
	Average maximum age (years)				
	Fecundity (eggs/spawning)				
	Average maximum size (cm)				
	Average size at maturity (cm)				
	Reproductive strategy				
	Mean trophic level				
		Average Productivity Score			
	Susceptibility Attribu	te Value	Score		
	Overlap of adult species range with fisher	ery			
	Distribution				
	Habitat				
	Depth range				
	Selectivity				
	Post-capture mortality				
		Average Susceptibility Score			
		PSA Risk Rating (From Table D3)			
	Compliance rating				
Standa	rd clauses 1.3.2.2				



## Table D2 - Productivity / Susceptibility attributes and scores.

Productivity attributes	Low productivity/ High risk	Medium productivity/ Medium risk	High productivity Low risk	
	Score 3	Score 2	Score 1	
Average age at maturity (years)	>4	2 to 4	<2	
Average maximum age (years)	>30	10 to 30	<10	
Fecundity (eggs/spawning)	<1 000	1 000 to 10 000	>10 000	
Average maximum size (cm)	>150	60 to 150	<60	
Average size at maturity (cm)	>150	30 to 150	<30	
Reproductive strategy	Live bearer, mouth brooder or significant parental investment	Demersal spawner "berried"	Broadcast spawner	
Mean trophic level	>3.25	2.5-3.25	<2.5	

Susceptibility attributes		High susceptibility/ High risk	Medium susceptibility/ Medium risk	Low susceptibility/ Low risk		
			Score 3	Score 2	Score 1	
Availability	Overlap of adult species range with fishery		>50% of stock occurs in the area fished	Between 25% and 50% of the stock occurs in the area fished	<25% of stock occurs in the area fished	
	2)	Distribution	Only in the country/ fishery	Limited range in the region	Throughout region/ global distribution	
Encounterability	1)	Habitat	Habitat preference of species make it highly likely to encounter trawl gear (e.g. demersal, muddy/sandy bottom)	Habitat preference of species make it moderately likely to encounter trawl gear (e.g. rocky bottom/reefs)	Depth or distribution of species make it unlikely to encounter trawl gear (e.g. epi-pelagic or meso-pelagic)	
	2)	Depth range	High overlap with trawl fishing gear (20 to 60 m depth)	Medium overlap with trawl fishing gear (10 to 20 m depth)	Low overlap with trawl fishing gear (0 to 10 m, >70 m depth)	
Selectivity			Species >2 times mesh size or up to 4 m length	Species 1 to 2 times mesh size or 4 to 5 m length	Species <mesh or<br="" size="">&gt;5 m length</mesh>	
Post capture mortality			Most dead or retained Trawl tow >3 hours	Alive after net hauled Trawl tow 0.5 to 3 hours	Released alive Trawl tow <0.5 hours	

**Note:** Availability 2 is only used when there is no information for Availability 1; the most conservative score between Encounterability 1 and 2 is used.



D3		Average Susceptibility Score			
		1 - 1.75	1.76 - 2.24	2.25 - 3	
Average Productivity	1 - 1.75	PASS	PASS	PASS	
Score	1.76 - 2.24	PASS	PASS	TABLE D4	
	2.25 - 3	PASS	TABLE D4	TABLE D4	