

IFFO RS Global Standard for Responsible Supply of Marine Ingredients

IFFO RS Limited

T: +44 (0) 2030 539 195 E: Standards@iffors.com W: www.iffors.com

Unit C, Printworks | 22 Amelia Street London, SE17 3BZ | United Kingdom





Global Standard for Responsible Supply of Marine Ingredients Fishery Assessment Methodology and Template Report V2.0



IFFO RS Global Standard for Responsible Supply of Marine Ingredients



Fishery Under Assessment	Squid Loligo vulgaris Northeast Atlantic
Date	July 2019
Assessor	Jim Daly

Application details and summary of the assessment outcome					
Name: Bioceval					
Address:					
Country: France		Zip:			
Tel. No.:		Fax. No.:			
Email address:		Applicant Code			
Key Contact:	Key Contact:		Title:		
Certification Body De	etails				
Name of Certification	n Body:	SAI Global Ltd	l		
Assessor Name	Peer Reviewer	Assessment Days	Initial/Surveillan approval	nce/Re-	Whole fish/ By- product
Jim Daly	Vito Romito	0.5	Initial		By-product
Assessment Period	2018				

Scope Details	
Management Authority (Country/State)	EU/Common Fisheries Policy
Main Species	Squid (Loligo vulgaris)
Fishery Location	North-East Atlantic
Gear Type(s)	Bycatch in trawls, some small-scale directed fisheries
Outcome of Assessment	
Overall Outcome	PASS
Clauses Failed	NONE
Peer Review Evaluation	Approve
Recommendation	Pass

Assessment Determination

Currently squid within the family *Loliginidae* are not assessed and there are no TAC constraints for these stocks. New data requirements have been launched to specific countries exploiting cephalopods in ICES areas. Data will be collated and advice published in due course.

IUCN has not yet categorised *Loligo vulgaris* (http://www.iucnredlist.org/search; accessed 16.02.18), nor does it appear in the CITES appendices (http://checklist.cites.org; accessed 04.07.19).

Using the Productivity-Susceptibility Analysis (PSA) for Category D species the assessor recommends that squid is approved under the current IFFO-RS Standard with a medium compliance rating due to selectivity and post-capture attributes.

Peer Review Comments

The Peer Reviewer agrees that because squid passes the PSA in category D, it should be approved as a byproduct under the current IFFO RS Standard.

Notes for On-site Auditor

Note: This table should be completed for whole fish assessments only.

Species-Specific Results

Category	Species	% landings	Outcome (Pass/Fail)
			A1
Cotogory			A2
Category A			A3
			A4
Category B			
Category C			
Category D	Squid Loligo vulgaris		PASS

[List all Category A and B species. List approximate total % age of landings which are Category C and D species; these do not need to be individually named here]

HOW TO COMPLETE THIS ASSESSMENT REPORT

This assessment template uses a modular approach to assessing fisheries against the IFFO RS standard.

Whole Fish

The process for completing the template for a **whole fish** assessment is as follows:

- 1. ALL ASSESSMENTS: Complete the Species Characterisation table, to determine which categories of species are present in the fishery.
- 2. ALL ASSESSMENTS: Complete clauses M1, M2, M3: Management.
- 3. IF THERE ARE CATEGORY A SPECIES IN THE FISHERY: Complete clauses A1, A2, A3, A4 for each Category A species.
- 4. IF THERE ARE CATEGORY B SPECIES IN THE FISHERY: Complete the Section B risk assessment for **each** Category B species.
- 5. IF THERE ARE CATEGORY C SPECIES IN THE FISHERY: Complete clause C1 for **each** Category C species.

- 6. IF THERE ARE CATEGORY D SPECIES IN THE FISHERY: Complete Section D.
- 7. ALL ASSESSMENTS: Complete clauses F1, F2, F3: Further Impacts.

A fishery must score a pass in **all applicable clauses** before approval may be recommended. To achieve a pass in a clause, the fishery/species must meet **all** of the minimum requirements.

By-products

The process for completing the template for **by-product raw material** is as follows:

- 1. ALL ASSESSMENTS: Complete the Species Characterisation table with the names of the by-product species and stocks under assessment. The '% landings' column can be left empty; all by-products are considered as Category C and D.
- 2. IF THERE ARE CATEGORY C BYPRODUCTS UNDER ASSESSMENT: Complete clause C1 for **each** Category C by-product.
- 3. IF THERE ARE CATEGORY D BYPRODUCTS UNDER ASSESSMENT: Complete Section D.
- 4. ALL OTHER SECTIONS CAN BE DELETED. Clauses M1 M3, F1 F3, and Sections A and B do not need to be completed for a by-product assessment.

By-product approval is awarded on a species-by-species basis. Each by-product species scoring a pass under the appropriate section may be approved against the IFFO RS Standard.

SPECIES CATEGORISATION

The following table should be completed as fully as the available information permits. Any species representing more than 0.1% of the annual catch should be listed, along with an estimate of the proportion of the catch each species represents. The species should then be divided into Type 1 and Type 2 as follows:

- **Type 1 Species** can be considered the 'target' or 'main' species in the fishery. They make up the bulk of annual landings and are subjected to a detailed assessment.
- **Type 2 Species** can be considered the 'bycatch' or 'minor' species in the fishery. They make up a small proportion of the annual landings and are subjected to relatively high-level assessment.

Type 1 Species must represent 95% of the total annual catch. Type 2 Species may represent a maximum of 5% of the annual catch (see Appendix B).

Species which make up less than 0.1% of landings do not need to be listed (NOTE: ETP species are considered separately). The table should be extended if more space is needed. Discarded species should be included when known.

The 'stock' column should be used to differentiate when there are multiple biological or management stocks of one species captured by the fishery. The 'management' column should be used to indicate whether there is an adequate management regime specifically aimed at the individual species/stock. In some cases it will be immediately clear whether there is a species-specific management regime in place (for example, if there is an annual TAC). In less clear circumstances, the rule of thumb should be that if the species meets the minimum requirements of clauses A1-A4, an adequate species-specific management regime is in place.

NOTE: If any species is categorised as Endangered or Critically Endangered on the IUCN Red List, or if it appears in the CITES appendices, it **cannot** be approved for use as an IFFO RS raw material. This applied to whole fish as well as by-products.

TYPE 1 SPECIES (Representing 95% of the catch or more)

Category A: Species-specific management regime in place. **Category B:** No species-specific management regime in place.

TYPE 2 SPECIES (Representing 5% OF THE CATCH OR LESS)

Category C: Species-specific management regime in place. **Category D:** No species-specific management regime in place.

Common name	Latin name	Stock	% of landings	Management	Category
Squid	Loligo vulgaris	North-East		No species specific	D
		Atlantic		management	
				regime	

CATEGORY C SPECIES

In a whole fish assessment, Category C species are those which make up less than 5% of landings, but which are subject to a species-specific management regime. In most cases this will be because they are a commercial target in a fishery other than the one under assessment. In a by-product assessment, Category C species are those which are subject to a species-specific management regime, and are usually targeted species in fisheries for human consumption.

Clause C1 should be completed for **each** Category C species. If there are no Category C species in the fishery under assessment, this section can be deleted. A Category C species does not meet the minimum requirements of clause C1 should be re-assessed as a Category D species.

Spec	cies N	ame			
C1	Categ	ory C Stock Status - Minimum Requirements			
\mathbf{v}	C1.1	Fishery removals of the species in the fishery under assessment are included in the			
		stock assessment process, OR are considered by scientific authorities to be negligible.			
	C1.2	The species is considered, in its most recent stock assessment, to have a biomass			
		above the limit reference point (or proxy), OR removals by the fishery under			
		assessment are considered by scientific authorities to be negligible.			
		Clause outcome:			
Evider	nce				
Doford	Deferences				
Kelele	References				
C 1	1 1	1222			
Standa	ira clau.	es 1.3.2.2			

CATEGORY D SPECIES

In a whole fish assessment, Category D species are those which make up less than 5% of landings and are not subject to a species-specific management regime. In the case of mixed trawl fisheries, Category D species may make up the majority of landings. In a by-product assessment, Category D species are those which are not subject to a species-specific management regime. In both cases, the comparative lack of scientific information on the status of the population of the species means that a risk-assessment style approach must be taken.

The process for assessing Category D species involves the use of a Productivity-Susceptibility Analysis (PSA) to further subdivide the species into 'Critical Risk', 'Major Risk' and 'Minor Risk' groups. If there are no Category D species in the fishery under assessment, this section can be deleted.

Productivity and susceptibility ratings are calculated using a process derived from the APFIC document "Regional Guidelines for the Management of Tropical Trawl Fisheries, which in turn was derived from papers by Patrick *et al* (2009) and Hobday *et al* (2007). Table D1 should be completed for each Category D species as follows:

- Firstly, the best available information should be used to fill in values for each productivity and susceptibility attribute.
- Table D2 should be used to convert each attribute value into a score between 1 and 3.
- The average score for productivity attributes and the average for susceptibility attributes should be calculated.
- Table D3 should be used to determine whether the species is required to meet the requirements of Table D4. A species which does not need to meet the requirements of D4 is automatically awarded a pass.
- Table D4 should be used to assess those species indicated by Table D3 to determine a pass/fail rating.
- Any Category D species which has been categorised by the IUCN Red List as Endangered or Critically Endangered, or which appears in the CITES appendices, automatically results in a fail.

1 Sp	ecies Name:	Squid Loligo vulgaris		
Pro	luctivity Attribute		Value	Score
Ave	rage age at maturity (ye	ars)	<1	1
Ave	rage maximum age (yea	rs)	1 - 1.5	1
Fecu	Indity (eggs/spawning)		1,000 - 10,000	1
Ave	rage maximum size (cm)	42	1
Ave	rage size at maturity (cr	n)	28	1
Rep	oductive strategy		Demersal	2
Mea	n trophic level		Large range of	2
	-		trophic levels*	Z
			Average Productivity Score	1.28
Susc	ceptibility Attribute		Value	Score
Ove	rlap of adult species ran	ge with fishery		
Dist	ribution		Throughout	1
			the region	1
Hab	itat		Neritic - semi	2
			pelagic	2
Dep	th range		Benthopelagic,	1
			0-500m	1
Sele	ctivity		Mesh size 70 -	3
			99 mm	
Post	-capture mortality		Long trawls	3
			Average Susceptibility Score	2
		PSA	Risk Rating (From Table D3)	PASS
			Compliance rating	MEDIUN

References:

R1: Arkhipkin et al (2015). World Squid Fisheries. Reviews in Fisheries Science & Aquaculture, vol. 23, pp 126 -133. <u>http://www.tandfonline.com/doi/pdf/10.1080/23308249.2015.1026226</u>

R2: Reference for mesh sizes. See Annex I page 19 for permitted mesh sizes when targeting squids (trawls). Regulation was amended in 2013 with no change on mesh size for this area:

Council Regulation (EC) 850/98 of 30 March 1998 for the conservation of fishery resources through technical measures for the protection of juveniles of marine organisms

http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:31998R0850&from=EN

R3: Guerra, A. & Rocha, F. (1994). The life history of Loligo vulgaris and Loligo forbesi (Cephalopoda: Loliginidae) in Galician waters (NW Spain). Fisheries Research, Volume 21, pp 43-69.

https://www.sciencedirect.com/science/article/pii/0165783694900957

R4: Sealifebase: http://www.sealifebase.org/summary/Loligo-vulgaris

(accessed 21.02.18)

R5* Reference notes: squids occupy a large range of range of trophic levels in marine food webs and show a large trophic width:

Coll et al (2013). Assessing the trophic position and ecological role of squids in marine ecosystems by means of food-web models: Deep Sea Research Part II: Topical Studies in Oceanography. Vol 95, pp 31-66. https://www.sciencedirect.com/science/article/pii/S0967064512001609

Standard clauses 1.3.2.2

Table D2 - Productivity / Susceptibility attributes and scores.

Productivity attributes	Low productivity/ High risk	Medium productivity/ Medium risk	High productivity/ Low risk	
	Score 3	Score 2	Score 1	
Average age at maturity (years)	>4	2 to 4	<2	
Average maximum age (years)	>30	10 to 30	<10	
Fecundity (eggs/spawning)	<1 000	1 000 to 10 000	>10 000	
Average maximum size (cm)	>150	60 to 150	<60	
Average size at maturity (cm)	>150	30 to 150	<30	
Reproductive strategy	Live bearer, mouth brooder or significant parental investment	Demersal spawner "berried"	Broadcast spawner	
Mean trophic level	>3.25	2.5-3.25	<2.5	

Susceptibility attributes		High susceptibility/ High risk	Medium susceptibility/ Medium risk	Low susceptibility/ Low risk	
		Score 3	Score 2	Score 1	
Availability	 Overlap of adult species range with fishery 	>50% of stock occurs in the area fished	Between 25% and 50% of the stock occurs in the area fished	<25% of stock occurs in the area fished	
	2) Distribution	Only in the country/ fishery	Limited range in the region	Throughout region/ global distribution	
Encounterability	1) Habitat	Habitat preference of species make it highly likely to encounter trawl gear (e.g. demersal, muddy/sandy bottom)	Habitat preference of species make it moderately likely to encounter trawl gear (e.g. rocky bottom/reefs)	Depth or distribution of species make it unlikely to encounter trawl gear (e.g. epi-pelagic or meso-pelagic)	
	2) Depth range	High overlap with trawl fishing gear (20 to 60 m depth)	Medium overlap with trawl fishing gear (10 to 20 m depth)	Low overlap with trawl fishing gear (0 to 10 m, >70 m depth)	
Selectivity		Species >2 times mesh size or up to 4 m length	Species 1 to 2 times mesh size or 4 to 5 m length	Species <mesh or<br="" size="">>5 m length</mesh>	
Post capture mortality		Most dead or retained Trawl tow >3 hours	Alive after net hauled Trawl tow 0.5 to 3 hours	Released alive Trawl tow <0.5 hours	

Note: Availability 2 is only used when there is no information for Availability 1; the most conservative score between Encounterability 1 and 2 is used.

D2		Average Susceptibility Score		
D3		1.00 - 1.75	1.76 - 2.24	2.25 - 3.00
Average Productivity	1.00 - 1.75	PASS	PASS	PASS
Score	1.76 – 2.24	PASS	PASS	TABLE D4
	2.25 - 3.00	PASS	TABLE D4	TABLE D4

D4	Spe	Species Name				
	Impa	cts On Species Catego	orised as Vulnerable by D1-D3 - Minimum Requirements			
	D4.1	D4.1 The potential impacts of the fishery on this species are considered during the management process, and reasonable measures are taken to minimise these impacts.				
	D4.2					
		·	Outcome:			
Evide	ciice					
Refer	ences					
Stand	ard claı	ıse 1.3.2.2				