



# MarinTrust Standard V2

# By-product Fishery Assessment VNM03 – Cod in ICES Subareas 1 & 2

#### **MarinTrust Programme**

Unit C, Printworks 22 Amelia Street London SE17 3BZ

E: standards@marin-trust.com

T: +44 2039 780 819



# Table 1 Application details and summary of the assessment outcome

	Species:	Cod (Gadus morhua)	
	Geographical area:	FAO 27 – Norwegian Sea and Barents Sea	
Fishery Under Assessment	Country of origin of the product:	Vietnam	
	Stock:	Cod in ICES Subareas 1 and 2 north of 67°N (Northern Norwegian coastal cod)	
Date	July 2023		
Report Code	VNM03		
Assessor		Sam Peacock	
Country of origin of the product - PASS	Vietnam		
Country of origin of the product - FAIL		n/a	

Application details and summary of the assessment outcome								
Company Name(s): Th	Company Name(s): Thien Quynh Co. Ltd, Thien Quynh Khanh Hoa Sole Member Limited Liability							
Company								
Country: Vietnam								
Email address:	Email address: Applicant Code:							
Certification Body Deta	ails							
Name of Certification E	Body:		LRQA					
Assessor Peer Reviewer		Assessment Days	Initial/Surveillance/ Re-approval					
Sam Peacock Jose Peiro Crespo 0.2 Surveillance 1								
Assessment Period		July 2023 -	- July 2024					

Scope Details	
Main Species	Cod (Gadus morhua)
Stock	Cod in ICES Subareas 1 and 2 north of 67°N (Northern Norwegian coastal cod)
Fishery Location	FAO 27 – Norwegian Sea and Barents Sea
Management Authority (Country/ State)	UK, EU, Norway
Gear Type(s)	Demersal trawls, gillnets, longlines
Outcome of Assessment	
Peer Review Evaluation	Pass
Recommendation	Pass



## Table 2. Assessment Determination

#### **Assessment Determination**

Cod has been categorised by the IUCN Red List as Vulnerable, and does not appear in the CITES appendices. Northern Norwegian coastal cod is managed relative to a reference point established in an international management plan, and was therefore assessed under Category C.

The most recent stock assessment was carried out in 2023 and utilised all commercial and recreational catch and bycatch data. The assessment concluded that stock biomass is above the reference point established by the management plan. For these reasons the byproduct continues to meet the MT requirements and should remain approved for use as a raw material.

#### **Fishery Assessment Peer Review Comments**

The by-product fishery under assessment is the Atlantic cod (*Gadus morhua*) demersal trawl, gillnet and longline fisheries in Norwegian Sea and Barents Sea ICES Subareas 1 and 2 north of 67°N (Northern Norwegian coastal cod) in FAO area 27. The species is classified as LC by the IUCN in European waters. The stock is managed relative to biomass-based reference points.

The stock was last assessed in 2023 by the ICES Arctic Fisheries Working Group (AFWG). According to that assessment, the stock biomass is above the reference point established by the management plan (SSB/owerbound). Therefore, the stock passes category C.

The peer review supports the auditor's recommendation to pass the Atlantic cod demersal trawl, gillnet and longline fisheries in Norwegian Sea and Barents Sea ICES Subareas 1 and 2 north of 67°N (Northern Norwegian coastal cod) under the Marin Trust IFFO RS v2.0 by-fishery standard for the production of fishmeal and fish oil.

Notes for On-site Auditor		
	 	 _



# **Species Categorisation**

**NB:** If any species is categorised as Endangered or Critically Endangered on the IUCN Red List, or if it appears in CITES Appendix 1, it **cannot** be approved for use as an MarinTrust raw material.

#### **IUCN Red list Category**

By-product material from a species listed by IUCN (the International Union for Conservation of Nature) under the Red List for the following categories shall immediately fail the assessment;

- EXTINCT (E) AND EXTINCT IN THE WILD (EW)
- CRITICALLY ENDANGERED (CR) facing an extremely high risk of extinction in the wild.
- ENDANGERED (EN) facing a very high risk of extinction in the wild.

By-product material may be used from the following categories provided that all clauses in the MarinTrust standard are passed.

- VULNERABLE (VU) facing a high risk of extinction in the wild.
- NEAR THREATENED (NT) does not qualify for above now, but is close or is likely to qualify for, a threatened category in the near future.
- LEAST CONCERN (LC) Widespread and abundant.
- DATA DEFICIENT (DD) and NOT EVALUATED (NE)

# Table 3 Species Categorisation Table

Common name	Latin name	Stock	Management	Category	IUCN Red List Category <sup>1</sup>	CITES Appendix 1 <sup>2</sup>
Cod	Gadus morhua	Northern Norwegian coastal cod	Yes	С	Vulnerable <sup>3</sup>	No

<sup>&</sup>lt;sup>1</sup> https://www.iucnredlist.org/

<sup>&</sup>lt;sup>2</sup> https://cites.org/eng/app/appendices.php

<sup>&</sup>lt;sup>3</sup> https://www.iucnredlist.org/species/8784/12931575



### **CATEGORY C SPECIES**

In a by-product assessment, Category C species are those which are subject to a species-specific management regime and are usually targeted species in fisheries for human consumption.

Clause C1 should be completed for each Category C species. If there are no Category C species in the fishery under assessment, this section can be deleted. Where a species fails this Clause, it should be assessed as a Category D species instead.

Spe	cies	Name	Cod			
<b>C1</b>	Category C Stock Status - Minimum Requirements					
CI	C1.1		ovals of the species in the fishery under assessment are included in the stock assessment are considered by scientific authorities to be negligible.	PASS		
	C1.2	reference po	is considered, in its most recent stock assessment, to have a biomass above the limit bint (or proxy), OR removals by the fishery under assessment are considered by scientific o be negligible.	PASS		
			Clause outcome:	PASS		


C1.1 Fishery removals of the species in the fishery under assessment are included in the stock assessment process, OR are considered by scientific authorities to be negligible.

Northern Norwegian coastal cod is subject to annual stock assessment by the ICES Arctic Fisheries Working Group (AFWG). The most recent assessment was an age-based analytical assessment conducted in 2023, which used catches in the model and forecast. The stock assessment included commercial and recreational landings data and bycatch, and ICES considers discarding to be negligible. The 2023 catch advice notes that due to changes in the tuning data, "the 2021 assessment results are...not directly comparable to the 2022-2023 results" (ICES 2023); however overall the results of the assessment are considered reliable. C1.1 is met.

C1.2 The species is considered, in its most recent stock assessment, to have a biomass above the limit reference point (or proxy), OR removals by the fishery under assessment are considered by scientific authorities to be negligible.

The 2023 catch advice includes an indication of the current status of the cod stock relative to reference points. Although there are no MSY or precautionary approach based reference points established for the stock, the reference point SSB<sub>lowerbound</sub> has been established by the management plan for the stock as "the limit above which the management plan is considered precautionary" (ICES 2023). The value for SSB<sub>lowerbound</sub> is 67,743t, and the most recent stock assessment estimated SSB in 2023 to be 74,654t. The catch advice states that "spawning-stock size is above SSB<sub>lowerbound</sub>" (ICES 2023). C1.2 is met.





Northern Norwegian coastal cod, estimated SBB relative to current reference point SSB<sub>lowerbound</sub> (ICES 2023)

#### References

ICES (2023). Cod (Gadus morhua) in subareas 1 and 2 north of 67°N (Norwegian Sea and Barents Sea), northern Norwegian coastal cod. In Report of the ICES Advisory Committee, 2023. ICES Advice 2023, cod.27.1-2coastN, https://doi.org/10.17895/ices.advice.21828306

Links	
MarinTrust Standard clause	1.3.2.2
FAO CCRF	7.5.3
GSSI	D.3.04, D5.01



# **CATEGORY D SPECIES**

Category D species are those which are not subject to a species-specific management regime. In the case of mixed trawl fisheries, Category D species may make up the majority of landings. The comparative lack of scientific information on the status of the population of the species means that a risk-assessment style approach must be taken.

D1	<b>Species Name</b>	n/a					
	Productivity Attribut	e Value	Score				
	Average age at maturity (years)						
	Average maximum age (years)						
	Fecundity (eggs/spawning)						
	Average maximum size (cm)						
	Average size at maturity (cm)						
	Reproductive strategy						
	Mean trophic level						
		Average Productivity Sco	re				
	Susceptibility Attribu	te Value	Score				
	Availability (area overlap)						
	Encounterability (the position of the s	• • • • • • • • • • • • • • • • • • •					
	within the water column relative to the	e fishing gear)					
	Selectivity of gear type						
	Post-capture mortality						
		Average Susceptibility Sco	re				
	PSA Risk Rating (From Table D3)						
	Compliance rating						
	Further justification for susceptibility scoring (where relevant)  For susceptibility attributes, please provide a brief rationale for scoring of parameters where there may be						
	uncertainty affecting your decision		,				
Refere	ences						
Stando	ard clauses 1.3.2.2						



# Table D2 - Productivity / Susceptibility attributes and scores.

Productivity attributes	High productivity (Low risk, score = 1)	Medium productivity (medium risk, score = 2)	Low productivity (high risk, score = 3)
Average age at maturity	<5 years	5-15 years	>15 years
Average maximum age	<10 years	10-25 years	>25 years
Fecundity	>20,000 eggs per year	100-20,000 eggs per year	<100 eggs per year
Average maximum size	<100 cm	100-300 cm	>300 cm
Average size at maturity	<40 cm	40-200 cm	>200 cm
Reproductive strategy	Broadcast spawner	Demersal egg layer	Live bearer
Mean Trophic Level	<2.75	2.75-3.25	>3.25

Susceptibility attributes			igh susceptibility igh risk, score = 3)				
Areal overlap (availability) Overlap of the fishing effort with the species range			10	10-30% overlap		>30% overlap	
Encounterability The position of the stock/species within the water column relative to the fishing gear, and the position of the stock/species within the habitat relative to the position of the gear	cosition of the description of t			fis en De	gh overlap with hing gear (high counterability). efault score for rget species		
Selectivity of gear type	а	Individuals < size at maturity are rarely caught	а	Individuals < size at maturity are regularly caught.	а	Individuals < size at maturity are frequently caught	
Potential of the gear to retain species	b	Individuals < size at maturity can escape or avoid gear.	Ь	Individuals < half the size at maturity can escape or avoid gear.	b	Individuals < half the size at maturity are retained by gear.	
Post-capture mortality (PCM) The chance that, if captured, a species would be released and that it would be in a condition permitting subsequent survival	re	ridence of majority eased post-capture d survival.	rel	idence of some eased post-capture d survival.	m	etained species or ajority dead when leased.	



D3		Average Susceptibility Score				
		1 - 1.75	1.76 - 2.24	2.25 - 3		
Average Productivity	1 - 1.75	PASS	PASS	PASS		
Score	1.76 - 2.24	PASS	PASS	TABLE D4		
	2.25 - 3	PASS	TABLE D4	TABLE D4		

<b>D4</b>	Spe	cies Name	n/a					
	Impac	ts On Species Categorise	ed as Vulnerable by D1-D3 - Minimum Requirements					
	<b>D4.1</b> The potential impacts of the fishery on this species are considered during the management							
		process, and reasonab	le measures are taken to minimise these impacts.					
	D4.2	There is no substantia species.	al evidence that the fishery has a significant negative impact on the					
			Outcome:					
Eviden	ice							
D4.2 T	here is r	no substantial evidence	that the fishery has a significant negative impact on the species.					
Refere	ences							
Links								
Marin <sup>*</sup>	Trust Sta	andard clause	1.3.2.2, 4.1.4					
FAO C	CRF		7.5.1					

D.5.01

GSSI